

# I.T.P


شبکه خبری و اطلاع رسانی صنعت مرغداری و دامپروری

شما فایل فوق را از سایت [www.ITPNews.com](http://www.ITPNews.com) دانلود کرده اید

**مالکیت مطلب و امتیاز آن برای این موسسه نیست**

لطفا در هنگام استفاده و یا برداشت مطلب و یا بخشی از آن، با ذکر نام نویسنده و یا صاحب امتیاز، تلاش نویسنده را بی ثمر نکنید

# BIRDS



Britannica Illustrated Science Library



## **About the pagination of this eBook**

Due to the unique page numbering scheme of this book, the electronic pagination of the eBook does not match the pagination of the printed version. To navigate the text, please use the electronic Table of Contents that appears alongside the eBook or the Search function.

For citation purposes, use the page numbers that appear in the text.

# BIRDS



**Britannica Illustrated Science Library**

Encyclopædia Britannica, Inc.

Chicago ■ London ■ New Delhi ■ Paris ■ Seoul ■ Sydney ■ Taipei ■ Tokyo



# Britannica Illustrated Science Library

© 2008 Editorial Sol 90

All rights reserved.

**Idea and Concept of This Work:** Editorial Sol 90

**Project Management:** Fabián Cassan

**Photo Credits:** Corbis, ESA, Getty Images, Daisy Gilardini—  
The Image Bank/Getty Images, Graphic News, NASA,  
National Geographic, Science Photo Library

**Illustrators:** Guido Arroyo, Pablo Aschei, Gustavo J. Caironi,  
Hernán Cañellas, Leonardo César, José Luis Corsetti, Vanina  
Fariás, Joana Garrido, Celina Hilbert, Isidro López, Diego  
Martín, Jorge Martínez, Marco Menco, Ala de Mosca, Diego  
Mourellos, Eduardo Pérez, Javier Pérez, Ariel Piroyansky, Ariel  
Roldán, Marcel Socías, Néstor Taylor, Trebol Animation, Juan  
Venegas, Coralia Vignau, 3DN, 3DOM studio

**Composition and Pre-press Services:** Editorial Sol 90

**Translation Services and Index:** Publication Services, Inc.

**Portions © 2008 Encyclopædia Britannica, Inc.**

Encyclopædia Britannica, Britannica, and the thistle logo are  
registered trademarks of Encyclopædia Britannica, Inc.

## Britannica Illustrated Science Library Staff

### Editorial

Michael Levy, *Executive Editor, Core Editorial*  
John Rafferty, *Associate Editor, Earth Sciences*  
William L. Hosch, *Associate Editor, Mathematics and  
Computers*  
Kara Rogers, *Associate Editor, Life Sciences*  
Rob Curley, *Senior Editor, Science and Technology*  
David Hayes, *Special Projects Editor*

### Art and Composition

Steven N. Kapusta, *Director*  
Carol A. Gaines, *Composition Supervisor*  
Christine McCabe, *Senior Illustrator*

### Media Acquisition

Kathy Nakamura, *Manager*

### Copy Department

Sylvia Wallace, *Director*  
Julian Ronning, *Supervisor*

### Information Management and Retrieval

Sheila Vasich, *Information Architect*

### Production Control

Marilyn L. Barton

### Manufacturing

Kim Gerber, *Director*

## Encyclopædia Britannica, Inc.

Jacob E. Safra, *Chairman of the Board*

Jorge Aguilar-Cauz, *President*

Michael Ross, *Senior Vice President, Corporate Development*

Dale H. Hoiberg, *Senior Vice President and Editor*

Marsha Mackenzie, *Director of Production*

International Standard Book Number (set):

978-1-59339-797-5

International Standard Book Number (volume):

978-1-59339-807-1

Britannica Illustrated Science Library: Birds 2008

Printed in China



[www.britannica.com](http://www.britannica.com)

# Birds





# Contents



## The Nature of Birds

Page 6



## The Art of Flying

Page 22



## The Lives of Birds

Page 40



## Diversity and Distribution

Page 62



## Humans and Birds

Page 80



# A Universe of Birds

**W**elcome to the world of birds. No matter how you approach it, this is a wonderful book not only for its pictures, splendid illustrations, size, and format but also because, as you read it, you will discover secrets about these inhabitants of the Earth, which, according to the history of evolution, came into being before humans. The text is written in a direct, easy-to-understand style. Most birds have a much-envied ability that has inspired poems and all types of experiments: they can fly. This enables them to see the Earth from afar, with its seas, mountains, rivers, cities, and other features. It has been estimated that more than 200 million birds migrate each year, all over the planet. Many of

**WHITE HERON** (*Egretta alba*)  
A species easy to distinguish in the proximity of rivers, lakes, and lagoons

them fly thousands of miles, crossing desolate deserts and windy seas to arrive in Africa or Antarctica. Some find their way using the sun, the moon, and the stars; others follow their parents or use the course of rivers or mountain chains as references. In general, smaller birds migrating across continents stop several times to get food. It is surprising how fast they travel, in spite of these stops: it has been calculated that some small species cover almost 2,500 miles (4,000 km) in five or six days. Several studies have shown that carrier pigeons and white-headed sparrows, for example, can travel more than 600 miles (1,000 km) per day. Some ducks, such as the blue-winged teal, complete their trip from Canada to central Mexico in about 35 days, making several stops to feed along the way.

**B**irds never cease to amaze us, whether hiding in trees, flying over high mountaintops, or nesting in Antarctica or on tall buildings. Perhaps the reason for such amazement is their behavior, which continues to be a mystery to human beings, as well as the differences among them. It is believed that there are approximately 9,700 living bird species in the world—more species than in any other vertebrate group except for fish. Once they reach adulthood, birds' weight varies from a

mere 0.06 ounce (1.6 g), in the case of hummingbirds, to as much as 330 pounds (150 kg) for African ostriches. Even though most birds fly, there are some—such as kiwis, rheas, and ostriches—that run quickly on the ground. Some birds, being perfectly adapted to aquatic life, live in oceans, rivers, and lakes. The shape of their feet and bills varies according to the environment in which they live. Some aquatic species have bills modified to filter small water particles, whereas birds of prey have strong bent bills to hold down and tear apart their prey. What is the diet of birds based on? Because of their great diversity and wide distribution, their diets differ greatly. In general, birds eat a bit of everything, although insects are the most important element of their diet. They eat fruit, seeds, nectar, pollen, leaves, carrion, and other vertebrates. Most birds lay their eggs in nests. Worthy of mention is the protective attitude that both males and females have toward their young. Adult birds care for their chicks, warn and protect them against the danger of predators, and guide them to safe places where they can live and feed. We invite you to investigate up close the world of these fascinating beings that are able to run, climb, swim, dive, and cross the skies. ●





# The Nature of Birds

**OWL** (*Bubo capensis*)  
This owl is native to Africa. It feeds on birds and mammals.

**BEYOND FEATHERS** 8-9

**ORIGIN** 10-11

**SKELETON AND MUSCULATURE** 12-13

**INTERNAL ORGANS** 14-15

**THE SENSES** 16-17

**DIFFERENT TYPES OF BILLS** 18-19

**EXPOSED LEGS** 20-21



**M**any scientists maintain that birds descended from dinosaurs because fossils of dinosaur specimens with feathers have been found.

As a group, birds have exceptional eyesight—they have the largest eyes in relation to the size of their bodies. In addition, they have very light bones, which are suitable for flight. Just like

their bills, birds' feet have also changed in accordance with the functions and particular needs of each species. For instance, walking birds—like other vertebrate groups—display a marked

tendency toward having a reduced number of toes; ostriches, for example, have only two. Some birds of prey, such as eagles, have feet that are veritable hooks. ●



# Beyond Feathers

Defining what a bird is brings to mind an animal covered with feathers that has a toothless bill and anterior extremities morphed into wings. Other distinguishing characteristics are that they are warm-blooded and have pneumatic bones—bones filled with air chambers instead of marrow. Birds have very efficient circulatory and respiratory systems and great neuromuscular and sensory coordination. ●




## Variety and Uniformity

We can find birds in every type of environment: aquatic, aerial, and terrestrial, in polar regions and in tropical zones. Their adaptation to the environment has been very successful. Nevertheless, birds are one of the groups that display the fewest differences among their members.

### WINE-THROATED HUMMINGBIRD

**0.06 ounce**  
(1.6 g)

WEIGHT OF THE SMALLEST BIRD



### AFRICAN OSTRICH

**330 pounds**  
(150 kg)

WEIGHT OF THE LARGEST BIRD

### PENGUIN

**-75 F**  
(-60° C)

THE TEMPERATURE PENGUINS CAN ENDURE IN ANTARCTICA

### WHITE-THROATED SPARROW

A small bird that lives in North America and on the Iberian Peninsula

## Adaptation to Flying

Some crucial anatomic and physiological characteristics explain birds' ability to fly. Their bodies and feathers reduce friction with the air and improve lift. Their strong muscles, light bones, air sacs, and closed double circulatory system also play a role in their ability to fly.

### FEATHERS

Unique. No other living animal has them. They are appealing for their structure, variety, and constant renewal.

### WINGS

propel, maintain, and guide birds during flight. They have modified bones and characteristic plumage.


**105.8° F**  
(41° C) IS THEIR BODY TEMPERATURE.

### FLIGHT FEATHERS

### UNDERTAIL COVERTS

### STRUCTURE

Balance in movement. A bird's internal architecture contributes to its stability. The location of its feet and wings helps to concentrate its weight close to its center of gravity.



### SENSES

Great visual acuteness and well-developed hearing

### INNER EAR

### EYE

### NAPE

### CREST


### NOSTRILS

### BILL

Originates in the epidermis. It is hard and resistant, with a consistency similar to that of horns. It grows continuously, like nails and feathers.

### IDENTIFICATION

There are differences in plumage and skin that make it possible to identify birds. The bill, because of its variations, also helps to establish bird groups.



### COVERTS

### THIGH

### TARSUS

### CHEST

### THORAX


### ABDOMEN

### High Metabolism

The high demands of flying are compensated by a high metabolic rate. Birds extract as many nutrients from food as they can.

### FEET

Birds walk on their toes. In general, they have three toes pointing forward and one pointing backward.



### TOES

### NAILS

### SONGBIRDS

Passeriformes, or passerines, form the most numerous group among birds; they are characterized by a well-developed syrinx that enables them to emit harmonious songs and trills and by a soft plumage of varied colors. Because of their brain development, it is believed that passerines were the most recent birds to come into existence.



# Origin

The evolution of birds is a debated theme in science. The most widespread theory states that birds descend from theropods, dinosaurs that walked on two legs. Fossils of dinosaur specimens with feathers have been found, but Archaeopteryx, a primitive bird that lived 150 million years ago, is the oldest relative known. Completely covered with feathers, it had a pair of wings that enabled it to fly. However, it retained many dinosaur traits.


## Archaeopteryx lithographica

lived in the Jurassic Period, 150 million years ago.





<b>Order</b>	Saurischians
<b>Suborder</b>	Theropods
<b>Diet</b>	Carnivore
<b>Length</b>	10 inches (25 cm)
<b>Height</b>	8 to 12 inches (20-30 cm)
<b>Weight</b>	18 ounces (500 g)

## From Reptile to Bird



## Fossils

Several fossil samples were found between 1861 and 1993. The first one, found in Bavaria, Germany, was very important because its discovery coincided with the publication of *On the Origin of Species* by Charles Darwin, at a time when the search for evolutionary "missing links" fascinated scientists. The original is located in the British Museum. Another fossil, which includes the head, is in the Berlin Museum.



**THREE TOES WITH TALONS**  
The hand has three extended fingers, each of which is equipped with a strong curved talon.

Talons for climbing trees





**WRIST**  
Its wrist joint was more flexible than that of modern birds, a trait it shared with dinosaurs.

**REPTILIAN JAWBONES WITH TEETH**  
Unlike modern birds, it did not have a horn bill. There was a tight row of sharp teeth on each jawbone.

**SPINE**  
Movable. The cervical vertebrae have a concave joint like that of the theropods, not a saddle-shaped one like that of birds.

**SKULL**  
Similar to that of present-day reptiles and early theropods. The arrangement of the brain and ears reveals that it had a great sense of orientation and that it was able to perform complicated maneuvers.



**FURCULA (Merged Collarbone)**  
Shaped like a boomerang, as in many theropods

**RIBS**  
Presence of ribs in the abdomen (gastralia), typical of reptiles and dinosaurs

**SAURIAN PELVIS**  
Hip and femur of the archosaurian, not avian, type

**UNMERGED METATARSUS**  
In modern birds, the tarsus and metatarsus are fused into the tarsometatarsus.

**VERTEBRATE TAIL**  
Composed of 21 or 22 pieces. Modern birds have tail vertebrae that are fused together into a single bone called the pygostyle.




During flight, it functioned as a rudder. On the ground, it provided balance for walking.

**TOES**  
The foot is functionally tridactyl. Its first toe (hallux), which usually points backward and typically does not touch the ground, is opposable, like that of modern birds (it can move in a direction perpendicular to toes II, III, and IV).

ARCHAEOPTERYX LITHOGRAPHICA  
Graphic Reconstruction

**FROM ARMS TO WINGS**  
It had a greater range of motion in the upper limbs than primitive dinosaurs.



ARCHAEOPTERYX  
150 million years ago

Its movements were limited by its shoulder joint, which was placed forward.

VELOCIRAPTOR  
99 to 65 million years ago

Birds have greater mobility than Archaeopteryx

PIGEON  
modern



# Skeleton and Musculature

Both lightweight and resistant, the skeleton of birds underwent important changes in order to adapt to flight. Some bones, like those of the skull and wings, fused to become lighter. Birds have fewer bones than other vertebrates. Because their bones are hollow, containing internal air chambers, the total weight of their bones is less than that of their feathers. Birds' spines tend to be very flexible in the cervical region and rigid near the rib cage, where a large, curved frontal bone called the sternum attaches. The sternum features a large keel, to which the pectoral muscles attach. These large, strong muscles are used for flapping the wings. In contrast, running birds, such as ostriches, have more developed muscles in their legs. ●


## Flapping Wings

Flying demands an enormous amount of energy and strength. Consequently, the muscles responsible for flapping the wings become very large, easily comprising 15 percent of the weight of a flying bird. Two pairs of pectorals, in which one muscle of the pair is bigger than the other, work to raise and lower the wings. They function symmetrically and in opposition to each other: when one contracts, the other relaxes. Their placement within the thoracic cavity corresponds roughly to the bird's center of gravity. The motion of the wings also requires strong tendons.




**HUMMINGBIRD**  
Because of its adaptation to stationary flight, its pectoral muscles can account for 40 percent of its total weight.

### DOWNWARD FLAP



### UPWARD FLAP



**SKULL**  
Light because of the fusing of bones, the skull does not have teeth, a bony jaw, or grinding muscles.

**UPPER MANDIBLE OF BILL**  
In some species, it is flexible.


**LOWER MANDIBLE OF BILL**  
It is flexible, allowing birds to open their mouths wide.

**FURCULA (COLLARBONE)**  
Known as the wishbone, it is unique to birds and results from the fusion of the collarbones.

**STERNUM**  
Hyperdeveloped in flying birds, the sternum's long keel facilitates the attachment of the pectorals.

### Wings

Without a doubt, wings are the greatest adaptation of birds. Strong tendons travel through the wings and merge into the hand bones, where the feathers are attached.



**THE COLOR OF THE FLESH**  
depends on the blood circulation in the muscles: the more circulation, the redder the flesh. Flying birds have red flesh, whereas nonflying birds, such as chickens, have white flesh.

**CERVICAL VERTEBRAE**  
Their number varies according to the type of bird. They make the neck flexible.

**CORACOIDS**

**HUMERUS**

**RADIUS**

**CARPAL BONES**


**ULNA**

**CARPOMETACARPUS**  
It is formed by the fusion of the hand bones.

**TOES**

**KNEE**


**FEMUR**




## Pneumatic Bones

Many of a bird's bones are pneumatic—that is, they are full of air instead of bone marrow. Some bones even have prolongations of air sacs. The bones may look fragile at first glance, but their incredible strength comes from a network of internal trabeculae (spongy bone structures), which resemble the trusses of a metal bridge.

**LEG MUSCLES**



**GRASPING DEVICE**  
When a bird is perched, it assumes a crouching position with its legs bent. This causes the tendons in its feet to tighten, which pulls its toes closed and locks its feet in place. This tendon-locking mechanism keeps birds from falling off branches while they sleep.



**SUPPORT POSITION**

**FEET**  
Birds have four toes, just like their ancestors, the reptiles.

**TARSOMETATARSUS**

**TOES**

**TIBIA**

**FALSE KNEE**

**PELVIS**

**PYGOSTYLE**  
The tail vertebrae are merged; the tail feathers are anchored to the tail.



# Internal Organs


Birds in flight can consume oxygen at a rate that a well-trained athlete would not be able to withstand for even a few minutes. Because of this oxygen consumption, all their organs have had to adapt. The lungs of birds, though smaller than those of mammals of similar size, are much more efficient. Their lungs have several air sacs that both increase the efficiency of their respiratory systems and make them lighter. A special feature of the digestive system is a crop in the esophagus, where food is stored for digestion or for feeding the young. A bird's heart can be four times larger in relation to its body size than a human's in relation to its body size. ●

## Digestive System

Birds have no teeth. They therefore ingest food without chewing, and their stomachs break it down. The stomach is divided into two parts: the glandular (or proventriculus) part, which secretes acids, and the muscular (or gizzard) part, whose muscular walls grind up what is eaten. In general, the process is very fast because flying requires a lot of energy, and the bird has to replenish that energy quickly. The digestive system ends at the cloaca, which is an excretory orifice shared with the urinary system. Birds absorb almost all the water they drink.

### FOOD ITINERARY

- STORAGE**  
Some birds have a crop, which enables them to store food and digest it later. This way they decrease their exposure to predators.
- PRODUCTION**  
The proventriculus secretes the gastric juices that initiate digestion.
- BREAKDOWN**  
In the gizzard, a strong and muscular pouch, food is broken down with the help of swallowed stones or sand. The stones and sand play the role of teeth.
- WATER ABSORPTION**  
occurs in the small intestine. Birds normally get water from the food they ingest.
- EXCRETION**  
The cloaca expels feces mixed with urine coming from the excretory system.



### TYPES OF GIZZARD


**Granivorous Birds** have thick muscle walls and strong mucous membranes (or internal skin) to break down seeds.



**Carnivorous Birds** have thin muscle walls because digestion takes place in the proventriculus.



- STOMACH
- LIVER
- GIZZARD
- PANCREAS
- SMALL INTESTINE




- TONGUE**  
Usually short, narrow, triangular, and not very muscular.
- ESOPHAGUS**
- TRACHEA**
- SYRINX**  
Makes it possible for birds to sing.
- CROP**
- LUNG**  
Almost rigid because of its structure.


## Respiratory System

Birds have the most efficient respiratory system of any vertebrate because of the great effort that flying demands. It has two small, almost rigid lungs that are assisted by nine air sacs distributed throughout the body. The air sacs work as bellows, but they do not carry out gas exchange. Oxygen enters the bloodstream through the parabronchi, which are much like the alveoli in human lungs, in that they serve as the tissue for gas exchange. In the parabronchi, blood and air flow past each other in tiny passages. Because air flows in one direction through the lungs, and blood in the lung capillaries flows in the opposite direction, birds can make use of all the air they inhale, much like fish can with their gills and in contrast with mammals, which cannot.

### THE AIR SACS



**SECTION OF THE LUNG**  
The reticulum formed by the parabronchi facilitates the exchange of gases with the blood.





**20%**

THE PERCENTAGE OF THE BODY'S VOLUME TAKEN UP BY LUNGS AND AIR SACS

A HUMMINGBIRD'S HEART BEATS **700** times a minute.

Rufous Hummingbird (*Selasphorus rufus*)




## A Highly Complex Heart

Similar to that of reptiles, but having a heart with four chambers instead of three, the circulatory system distributes nutrients and oxygen throughout the body according to the body's needs. The heart's size and rate vary, depending on the bird's weight and activities. In general, bigger birds have smaller and slower hearts. For example, the heart of a seagull on the ground beats 130 times a minute; in flight, it beats 625 times a minute. A hummingbird's heart can beat 700 times a minute.

### THE HEART'S ASYMMETRY

The left side of the heart is more developed, because it pumps blood to the whole body. The right side pumps blood only to the lungs.



- The Blood enters** through the right and left arteries.
- Relaxed Ventricles**  
They open the atrioventricular valves.
- Contracted Ventricles**  
The blood enters the bloodstream.





# The Senses

In birds, the sense organs are concentrated on the head, except for the sense of touch, which is found all over the body. Birds have the largest eyes with respect to the size of their bodies. This enables them to see distant objects with considerable precision. Their field of vision is very broad, over 300 degrees, but in general they have little binocular vision. The ear—a simple orifice, but very refined in nocturnal hunters—helps them notice sounds inaudible to humans, which facilitates the detection of prey while flying. The senses of touch and smell, on the other hand, are important only to some birds, and the sense of taste is almost nonexistent. ●

## The Ear

Birds' ears are simpler than those of mammals: a bird's ear has no outer portion, and in some cases it is covered with rigid feathers. A notable part of the ear is the columella—a bone that birds share with reptiles. The ear is nonetheless well developed, and birds have very acute hearing; whereas human beings can detect just one note, birds can detect many. The ear is essential to a bird's balance, a key factor in flying. It is also believed that in certain species the ear works as a barometer, indicating altitude.



### LOCATION OF THE EARS

Located at different heights on the head, the ears cause the sense of hearing to occur with a slight delay. In nocturnal hunters, such as owls, this asymmetry allows for the triangulation of sounds and the tracking of prey with a minimal margin of error.

## Touch, Taste, and Smell


The sense of touch is well developed in the bill and tongue of many birds, especially in those birds that use them to find food, such as shore birds and woodpeckers. Usually the tongue is narrow, with few taste buds, but they are sufficient to distinguish among salty, sweet, bitter, and acidic tastes. The sense of smell is not very developed: although the cavity is broad, the olfactory epithelium is reduced. In some birds, such as kiwis and scavengers (condors, for example), the olfactory epithelium is more developed.



## Vision


is the most developed sense in birds because some flight maneuvers, as well as the recognition of food from afar, depend on it. Birds have relatively large eyes. In most cases, they are wider than they are deep because the lens and the cornea—which is supported by a series of sclerotic bony plates—project beyond the eye socket. In hunting birds, the eyes are almost tubular.

The muscles around the eye change its shape, alter the lens, and create greater visual acuity: birds typically have a 20-fold magnification (and sometimes, as in the case of some diving birds, a 60-fold magnification), in comparison with humans. Their sensitivity to light is also remarkable, with some species being able to recognize light spectra invisible to the human eye.




### FIELD OF VISION

The eyes—when located on the sides of the head, as is the case with most birds—create a broad field of vision: more than 300 degrees. Each eye covers different areas, focusing on the same object only when looking ahead through a narrow binocular field of vision.



### THE HUMAN FIELD OF VISION

The eyes, located at the front, move together, covering the same area. Because human beings cannot move their eyes independently from each other, they have only binocular vision.




### COMPARISON OF BINOCULAR FIELDS OF VISION


Binocular vision is essential for measuring distances without making mistakes. The brain processes the images that each eye generates separately as if they were a single image. The small differences between the two images

allow the brain to create a third one in depth, or in three dimensions. Hunting birds, for which the correct perception of distance is a life-and-death matter, tend to have eyes located toward the front, with a wide field of binocular vision.

In contrast, birds with lateral eyes calculate distance by moving their heads, but they record a larger total field of vision to avoid becoming prey. Owls are the birds with the greatest binocular vision—up to 70 degrees.



**HUNTING BIRDS' FIELD OF VISION**  
Frontal eyes reduce the total field of vision but allow for a wide field of binocular vision.




**NONHUNTING BIRDS' FIELD OF VISION**  
The lateral eyes open the field of vision to as much as 360 degrees but reduce the binocular field.





# Different Types of Bills

The beak, or bill, is a projecting structure of horn—made out of the same material as the nails—that grows as it is worn down. In the case of adult birds, bill size remains constant. The bill is joined to the skull in a way that allows for the movement of the lower mandible and, thus, the opening of the mouth. Most birds depend on their bills to get food. There are many types of bills, which differ in size, shape, color, and hardness, depending on the way in which the bird gets its food. ●



## COMPOSITION AND STRUCTURE

The jaws are covered with a hard horn layer called the ramphotheca, which is the external, visible portion. This determines the bill's color.

## Parts of the Bill


Each jaw has characteristic elements. In the upper one, from the back to the front, are the nostrils (or nasal cavities), the culmen (or maxillary cover), and the tip, which, in carnivorous birds, contains the tomial, or killing, tooth. In the lower jaw is the gonyx, or cover. The variations found in each part of the bill are conditioned by the bill's function.

## Hardness

Its long, stout bill is extraordinarily hard. Despite its appearance, the bill is very light, and birds can use it adeptly to seize and to open the fruits they eat.

## Heterogeneous Shapes

Bills have a wide array of names and shapes, but they are usually classified according to their length in relation to the head (short or long); to the curvature of its axis (pointing upward or downward); to its width; to its general shape (conical, stiletto-shaped, or spatula-shaped); and to the presence or absence of accessory pieces, such as grooves, horny plates, or false serrated teeth.



**TOUCANS AND ARICARI**  
With their long, thick bills, they can reach fruit located on branches that are too thin for the bird to sit on. Their bills are also used to break the peels and seeds of fruits.



**FLAMINGO**  
Flamingos have thin, threadlike structures inside their bills whose function is similar to that of the baleen of whales. They feed on microorganisms through filtration.



**HERON**  
The heron fishes in shallow waters and has a long, solid, sharp bill that quickly slices through the water to easily harpoon fish.



**GREENFINCH**  
Like granivores in general, it has a strong, conical bill, used to detach seeds from plants and, sometimes, to crack them.



**FALCON**  
It uses the false (tomial) tooth at the tip to detach the flesh from the bone and to break the spine of its prey.



**RAVEN**  
Because of its unrestricted diet, its bill is simple and relatively long.



**HUMMINGBIRD**  
The ability to reach the bottom of a flower in order to suck nectar requires not only a long, thin bill but also a special tongue.



**CROSSBILL**  
It feeds only on pine seeds. It uses its bill to reach the scales of pine cones, open them, and extract the pine nuts.

## You Are What You Eat

There is a close relationship between a bird's bill and its diet. Because the bill serves to pick up, hunt, tear, and transport the food, depending on the bird's lifestyle, its appearance is related to the bird's diet. If the

diet is very specific, the bill tends to have an adapted, unique shape, as is the case with hummingbirds. Omnivorous birds, on the other hand, have simple bills with no special alterations that are suitable for all tasks.






# Exposed Legs

Taking a quick look at the extremities of birds, including their toes and claws, can help us learn about their behavior. The skin of their legs and feet can have some striking features. All these characteristics reveal information about the environments in which different groups of birds live, as well as about their diets. Scientists use these characteristics as a basis for classifying birds. The detailed study of the anatomy of a bird's leg and foot can offer useful information. The shape and placement of bones, muscles, and tendons make it possible to understand how birds hold their prey or perch on branches, as well as to learn about the mechanics of their movement across the ground and in the water. ●

## Different Types

The foot usually has four toes. Three of them have a similar size and position. Opposite them is a smaller toe called the hallux. This pattern varies among different bird groups. For example, the position and shape of toes can differ. There are even cases in which two toes are functional while the others have been reduced in size. This is the case with flightless birds such as rheas. Differences are also found in the skin, which may form a web between the toes and projections of horn. All these characteristics become tools to help the bird survive in its environment and face challenges regarding obtaining food.

**FEET DESIGNED FOR SEIZING**  
Found on birds of prey and nocturnal rapacious birds. Their feet are strong, and their toes end in long, curved, sharp claws. They seize prey and transport it in flight.



**FEET DESIGNED FOR CLIMBING**  
Found on parrots, woodpeckers, and cuckoos. The hallux and the fourth toe are pointed backward. This arrangement provides the birds with more strength for climbing tree trunks.




**FEET DESIGNED FOR WALKING**  
Found on herons, flamingos, and storks. The toes and legs are very long. The hallux is pointed backward. They live in places with soft ground, such as swamps and river banks.




**FEET DESIGNED FOR PERCHING**  
Found on hummingbirds, kingfishers, ovenbirds, and nightjars. They have small feet, with the second, third, and fourth toes joined together. This makes it possible for them to stand still.



**FEET DESIGNED FOR SWIMMING**  
Found on auks, ducks, and penguins, which have a membrane between their toes that forms a web and increases the surface of the foot that is in contact with the water.



**FEET DESIGNED FOR RUNNING**  
Found on bustards, curlews, and rheas. They have long legs with short toes. The hallux and the fourth toe are very small, which decreases contact with the ground while running.



**THE FOOT II**  
The distal tarsal bones merge into the metatarsal bone and create tarsometatarsal bones.

**THE FOOT I**  
Toes 1 (hallux) and 2 have three phalanges, toe 3 has four, and toe 4 has five.



**TRICOLORED HERON**  
Its feet have long, thin toes that allow it to move on soft ground, such as in swamps, on river banks, and on lake shores. It lives in the regions of Arica and Coquimbo in Chile.

**Adaptation to Trees**  
The common waxbill perches and sleeps on tree branches without expending much energy. The weight of the body alone causes its toes to close tightly around the branch.

## Claws, Scales, and Spurs

These striking foot structures play a role in finding food, movement, protection, and defense, among other things. The claws can be long and sharp, in the case of birds of prey, or short and round, in the case of walking birds. Owls have a comblike claw that they use to groom their plumage. Their scales, inherited from reptiles, help protect their feet. In some cases, they help the birds to move through water. Many birds, such as chickens, pheasants, and crested screamers (a South American waterbird), have a spur, which they use as a defensive or offensive weapon.




**BALD EAGLE (Talons)**  
Very long, curved, pointed claws. They envelop the body of the prey and pierce it.

**KNEE AND THIGH**  
The thigh is included in the body and has a shortened femur. The knee is near the center of gravity.

**TIBIA**  
The tibia merges into tarsal bones and forms the tibiotarsus. It has a slightly developed fibula on its lateral face.



**GREAT CRESTED GREBE (Lobed Toes)**  
In some swimming birds, the toes look like oars. They have a continuous wide border.



## Internal/External Structure

Birds walk on their toes, which form the first portion of their feet. The second portion is formed by the tarsometatarsus. Its top part is connected to the tibia, through a joint similar to that of our ankle. That is why the leg flexes backward. The knee, equivalent to

ours, is higher up and works like a hip. It is located close to the body, and it helps to maintain balance. The thigh bone also stabilizes the body by adding weight to the skeleton. All the movements of these bones are controlled by tendons and muscles.



**SCOTS DUMPY ROOSTER (Spurs)**  
The spurs originate in skin and bone tissues. When males fight over territory or over a female, they use their spurs to defend themselves.



# The Art of Flying

**PARROT FEATHER**  
Detail of the feathers worn by  
these colorful aerial acrobats

**ADAPTATIONS 24-25**

**FEATHERS 26-27**

**WINGS TO FLY 28-29**

**TAIL TYPES 30-31**

**TO RENEW IS TO LIVE 32-33**

**GLIDING 34-35**

**FLAPPING FLIGHT 36-37**

**SPEED RECORDS 38-39**



**B**irds move in the air the same way a glider does, that is, by making the most of air currents to gain height and speed while moving. The

shape of the wings varies according to the needs of each bird group. Some cover considerable distances and thus have long, narrow wings, whereas others have short, rounded

wings that allow them to make short flights from branch to branch. Birds also have shiny, colorful feathers that males frequently use both to attract females and to hide from enemies.

Feathers are usually renewed once a year, and this process is as vital to birds as feeding. ●



# Adaptations

There are three main theories to explain why birds developed the ability to fly. The evidence that supports each of them tells a story of adaptations to an aerial world in which the fight for food and survival is key. One reasonable theory argues that birds descended from an extinct line of biped reptiles that fed on plants and used to jump from branch to branch to flee. ●


## From Reptile to Bird

It is known that several evolutionary lineages from both reptiles and birds did not survive the evolutionary process, and that the lineage that truly links these two animal groups has not yet been found. However, some theories state that the change from reptile to bird took place through a long process of

adaptation. There are two arguments and a variant: the arboreal theory, which posits an air-ground flight model; the cursorial (or running) theory, which focuses on the need for stability when running; and a variant, related to parental care, which posits that dinosaurs started to fly as a way of keeping their eggs safe.

### 1 THE ARBOREAL THEORY

This theory, the most accepted for a long time, states that flight was an adaptation to the environment in which certain herbivorous climbing reptiles lived. At first, dinosaurs developed a kind of parachute to protect them if they missed a branch when jumping, and later it became a way to move from tree to tree. Finally, flight evolved to involve the flapping of wings, which allowed the animal to cover greater distances.



### 2 THE CURSORY, OR RUNNING, THEORY

Supported by good fossil evidence, the running theory argues that birds descended from certain bipedal dinosaurs that were fast runners. Their arms opened, evolving into wings, to stabilize them as they jumped. Progression from this development to flying was simply a matter of time.



### 3 PARENTAL CARE VARIANT

This variant proposes that reptiles started to climb trees to prevent their young from becoming prey. Gliding removed the need to climb out of the trees.




## Today

The tips of the wings propel flight; the arms support the bird; and the shoulders enable the flapping movements.


150 MILLION YEARS AGO

The bones are extended and reinforced; then they merge.



175 MILLION YEARS AGO

The shoulders can perform a wider range of movements. The fingers merge.



200 MILLION YEARS AGO

Dinosaur arm with pincer claw and limited movement



### THE EMERGENCE OF THE WING

It evolved from an arm with a talon into a limb, without a talon, that was adapted for flight. The causes of this change are not yet clear to scientists. However, fossil records show how bones merged until they reached their present forms.

### EAGLE

In its maneuvers, this great hunter displays the entire evolution of flight.

### OTHER FLYING ANIMALS

In flying animals, from primitive pterodactyls to bats, wings have always been a flap of skin. A tear creates serious problems because it takes time to heal, and the wing may be misshapen afterward.

### GLIDING SPECIES



### THE BEST SOLUTION

Feathers are a unique evolutionary advantage. Their versatility, strength, individual nature, and ease of replacement make them an ideal adaptation to flight for vertebrates.

### 4 FEATHERS

Today found only on birds, feathers are scales partitioned into three smaller sections. They form a light, uniform, resistant network that covers the whole body.

26 pounds (12 kg)

IS THE MAXIMUM WEIGHT AN EAGLE CAN CARRY DURING FLIGHT.

The eagle itself usually weighs about 13 pounds (6 kg) and can generally carry prey weighing 6.5 pounds (3 kg). Some eagles, however, have been seen carrying prey estimated to weigh 13 pounds (6 kg). Carrying any more weight would require bigger wings, which would be more difficult to move and less efficient. It is believed that large flying animals disappeared because of this limitation.



### FROM SCALES TO FEATHERS

The development of feathers brought great advantages to birds because feathers enabled them to fly. Feathers evolved from scales, and they are made of the same material. Feathers keep the body's temperature constant and are lighter than scales.




# Feathers

Feathers are the feature that distinguishes birds from all other animals. They make birds strikingly colorful, protect them against cold and intense heat, enable them to move easily through the air and water, and hide them from enemies. Feathers are also one of the reasons why human beings have domesticated, caught, and hunted birds. A bird's set of feathers is called its plumage, and its color is essential for reproductive success. ●



## WHAT IS KERATIN?

Keratin is a protein that forms part of the outermost layer of a bird's skin, just as it does in other vertebrate animal groups. Keratin is the main component of feathers, hair, and scales. Its distinct resistance helps keep the hooklets woven together in the vane. This allows birds' feathers to maintain their shape in spite of the pressure exerted by the air during flight.



25,000


THE NUMBER OF FEATHERS THAT LARGE BIRDS, SUCH AS SWANS, CAN HAVE. In contrast, the number of feathers small birds, such as songbirds, can have varies between 2,000 and 4,000.



TRAILING EDGE NOTCH The turbulence during flight is reduced by this notch, found near the tip of the wing.

## Structure

The structure of feathers has two parts: a shaft and a blade. The shaft is called the rachis, and the part connected to the bird's skin is called the calamus. The movement of a feather is generated in the rachis. The blade is composed of barbs that branch into barbules. The feather's blade, in which the barbules have a series of barbicels, or hooklets, at the tip, is called a vane. The interlocking hooklets in the vane create a network that adds rigidity and resistance to the feather. It also defines the characteristic aerodynamic shape of feathers and helps make the feather waterproof. When feathers wear out, birds have the ability to replace them with new ones.



## INFERIOR UMBILICUS

The orifice at the base of the calamus, into which the dermic papilla penetrates. New feathers receive nourishment through it.

HOLLOW INTERIOR

INNER PULP OF THE SHAFT

## CALAMUS

It provides the necessary nutrients for feathers to grow. Nerve endings that stimulate the feather's movement are found at its base. This allows the bird to detect changes in its surroundings.

## SUPERIOR UMBILICUS

It contains some loose barbs. Some feathers have a secondary rachis, the hyporachis.

## BARBS

are slim, straight ramifications that grow perpendicular to the rachis.

## EDGE

The edge presents an excellent aerodynamic profile for flying.

## RACHIS

A feather's main shaft, similar to a hollow rod

## VANE, OR BLADE

Its outer portion contains a great number of barbicels.

## PREENING THE PLUMAGE

Birds need to preen their feathers with their bills not only to keep them clean and free of parasites but also to keep them lubricated, which helps birds resist inclement weather. Birds touch their uropygial, or preen, glands with their bills. Then they distribute the oil and wax this gland produces all over their plumage. This task is a matter of survival.



## SELF-CLEANING WITH ANTS

Some birds, such as certain tanagers, catch ants with their bills and grind them. They then oil their feathers with the ground-up ants. It is believed that the acid juices from the squashed ants work as a repellent against lice and other external parasites.

## DUST BATH


Birds such as pheasants, partridges, ostriches, pigeons, and sparrows perform dust baths to control the amount of grease on their feathers.

## Types of Feathers

There are three main types of feathers, classified according to placement: those closest to the body are down, or underlying feathers; those at the top are contour feathers; and those on the wings and tail are flight feathers, which are often referred to as remiges (on the wings) and rectrices (on the tail).

## DOWN

These light and silky feathers protect the bird against the cold. They have a short rachis, or none at all. Their barbs are long, and their barbules lack hooklets. In general, down is the first type of feather that birds develop when they hatch.




## CONTOUR

Also called covert feathers, they are short and rounded. They are more rigid than down feathers. Because they cover the body, wings, and tail, they give birds their shape as they fly.




## SPECIAL FEATHERS

Vibrissae are special feathers formed by only one filament. Sometimes they have loose barbs at the base that perform a tactile function. They are located at the base of bills or nostrils or around the eyes. They are very thin and are usually blended with contour feathers.



## PTERYLAE AND APTERIA

At first glance, a bird's body is covered with feathers. However, feathers do not grow all over the body but rather in particular areas called pterylae. This is where the papillae, which create new feathers, are found. The shape and placement of pterylae vary according to species. Pterylae are surrounded by naked areas, called apteria, in which feathers do not grow. Penguins are the only birds whose bodies are completely covered with feathers. This characteristic makes it possible for them to live in cold regions.



## IMPERIAL HERON

Powder down keeps its plumage waterproof.

## POWDER DOWN

This special type of feather can be found on some aquatic birds. They grow constantly and break off at the tip into small waxy scales. This "powder" is preened into the plumage to provide protection.




# Wings to Fly

**W**ings are highly modified arms that, through their unique structure and shape, enable most birds to fly. There are many types of wings; they vary by species. For instance, penguins, which are flightless, use their wings for the specialized task of swimming. Among all wings that have existed in the animal kingdom, those of birds are the best for flying. Their wings are light and durable, and in some cases their shape and effectiveness can be modified during flight. To understand the relationship between wings and a bird's weight, the concept of wing loading, which helps explain the type of flight for each species, is useful. ●

## Wings in the Animal Kingdom

Wings have always been modified arms, from the first models on pterosaurs to those on modern birds. Wings have evolved, beginning with the adaptation of bones. Non-avian wings have a membranous surface composed of flexible skin. They extend from the bones of the hand and body usually down to the legs, depending on the species. Avian wings, on the other hand, are based on a very different principle: the arm and hand form a complex of skin, bone, and muscle, with a wing surface consisting of feathers. Furthermore, the avian wing allows for important changes in form, depending on the bird's adaptation to the environment.




**PTERODACTYLS**  
still had talons, and only one finger extended their wings.

**BATS**  
Four fingers extend the membrane, and the thumb remained as a talon.

**BIRDS**  
The fused fingers form the tip of the wing where the rectrices, or primary feathers, are attached.

## Wing Size and Loading

The wingspan is the distance between the tips of the wings. Together with width, it determines the surface area, which is an essential measurement for bird flight. Not just any wing can support any bird. There is a close relationship between the animal's size (measured by weight) and the surface area of its wings. This relationship is called wing loading, and it is crucial in understanding the flight of certain species. Albatrosses, with large wings, have low wing loading, which makes them great gliders, whereas hummingbirds have to flap their small wings intensely to support their own weight. The smaller the wing loading, the more a bird can glide; the bigger, the faster a bird can fly.



**ARGENTAVIS MAGNIFICENS** (extinct)

**LARGER FINGER**  
**SMALLER FINGER**  
**CARPOMETACARPUS**  
**ALULAR DIGIT**  
Controls the alula, a feathered projection on the front edge of the wing.  
**ULNA**

**RADIUS**

**HUMERUS**

**CORACOID**

**STERNUM OR KEEL**

**PRIMARYS**  
They are in charge of propulsion; they are also called remiges.

**PRIMARY COVERTS**  
They cover the remiges and, with the alula, change the wing shape at will.

**MEDIAN WING COVERTS**  
They change the wing's lift when they rise slightly.

**GREATER WING COVERTS**  
They create more surface area and cover the intersection point of the tertiaries.

**SECONDARIES**  
Their number varies greatly depending on the species. They complete the surface.

**TERTIARIES**  
Together with the secondaries, they create the wing's surface.

**LOOSE FEATHERS**  
Sometimes barbels are missing, and feathers on the wing come apart, creating a loose and ruffled appearance.

**PRIMARY FEATHERS**  
Flying birds have from nine to 12 primary feathers. Running birds may have up to 16.

**FUNCTION**  
The wings of ostriches carry out the functions of balancing, temperature regulation, and courtship.


## Types of Wings

According to the environment in which they live and the type of flight they perform, birds have different wing shapes that allow them to save energy and to perform efficiently during flight. The wing shape also depends on the bird's size. Consequently, the number of primary and secondary feathers changes depending on the needs of a given species.

<p><b>The external primary feathers are longer.</b></p> <p><b>FAST WING</b> Remiges are large and tight to allow for flapping; the surface is reduced to prevent excessive friction.</p>	<p><b>The outermost primary feathers are shorter than the central ones.</b></p> <p><b>ELLIPTICAL WINGS</b> Functional for mixed flights, they are very maneuverable. Many birds have them.</p>	<p><b>They are wide at the base, with separate feather tips.</b></p> <p><b>WINGS FOR SOARING ABOVE LAND</b> Wide, they are used to fly at low speeds. The separate remiges prevent turbulence when gliding.</p>	<p><b>There are many secondary feathers.</b></p> <p><b>WINGS FOR SOARING ABOVE THE OCEAN</b> Their great length and small width make them ideal for gliding against the wind, as flying requires.</p>	<p><b>Short feathers are located all over the wing.</b></p> <p><b>WINGS FOR SWIMMING</b> In adapting to swimming, the feathers of penguins became short, and they serve primarily as insulation.</p>
--	--	---	---	--

## Flightless Wings

Among these, penguins' wings are an extreme case of adaptation: designed for rowing underwater, they work as fins. On running birds, wings' first and foremost function is to provide balance as the bird runs. These wings are also related to courtship, as birds show off their ornamental feathers during mating season by opening their wings or flapping them. Wings are also very efficient at controlling temperature, as birds use them as fans to ventilate their bodies.







Over the course of evolution, birds' tail vertebrae fused into a pygostyle, and in their place feathers of different sizes and colors emerged. These feathers have multiple uses: they can control aerial maneuvers during flight, work as brakes during landing, and make noise. Males also use them during courtship to dazzle and win over females. Usually the tail is formed by rectrices that vary in number, length, and rigidity depending on the species. ●

## Courtship Display

▶ The tail feathers of the female black grouse are straight, whereas those of the male have a half-moon shape. They usually keep the feathers closed and near the ground, but during the courtship displays they spread them out and show them off completely. To finish the show, the male runs back and forth in front of the female.



**RECTRICES**  
Tail feathers can wear out and fray because of friction during flight or by brushing against vegetation.



**UNDERTAIL COVERTS**  
Feathers that cover the lower part of rectrices, protecting them against the wear and tear caused by air friction.

**Black Grouse**  
*Lyrurus tetrix*  
The male is recognized by its bluish black plumage and the red caruncle over its eyes.

## Fan of Rectrices

▶ On flying birds, it is light and aerodynamic. On tree-climbing birds, such as woodpeckers, the plumage is rigid, which allows them to use it as a support (pointed tail). The coverts of male peacocks are more developed than their rectrices so that the peacock can show them off.

**FORKED TAIL**  
Found on swallows and frigate birds. The external feathers are very long and look like scissors.



**ROUNDED TAIL**  
Found on some songbirds. The central feathers are only slightly longer than the external ones.



**GRADUATED TAIL**

Found on trogons and kingfishers. When closed, the tail has a layered shape.



**MARGINATED TAIL**  
Found on blue jays. The central feathers are only slightly shorter than the external ones.



**SQUARE TAIL**  
Found on quails. The tail is short, with even-sized feathers.



## The Key to How It Works

▶ The tail can perform a variety of functions because of the movement and shape of the feathers. The powerful muscles in the pygostyle prepare the plumage for courtship displays and for flight, provide balance in walking and alighting on trees, and work as rudders for swimming.

**OPEN**



**LANDING I**  
The plumage spreads out, and the main axis of the body is positioned parallel to the ground.

**CLOSED**



**LANDING II**  
The body leans backward, and the tail closes. The legs prepare to grab the branch.

**OPEN**



**LANDING III**  
The spread-out tail feathers, together with an intense flapping of the wings, make it possible for the bird to slow down and prepare its body to land.





# To Renew Is to Live

The periodic renewal of plumage is called molting. It is the replacement of worn-out, older feathers with new ones that are in better condition. In a bird's life cycle, molting is as important an event as migrating or caring for young. The beginning of this phenomenon is determined by environmental factors that trigger a series of hormonal stimuli in birds: they start to eat more and to decrease their other activities. This, in turn, causes them to gain weight through an accumulation of fat that will serve as the source of energy for developing new plumage. ●

## Plumage Molting

The main function of molting is to replace worn-out plumage. It also helps the bird adapt its appearance to the seasons and to different stages in life. The renewal can be partial or total. Some feathers are replaced before the spring, when the task is to attract a partner for reproductive purposes. In the fall, before birds have to start caring for their young, the renewal is complete. On most birds, molting takes place in each pterylya, following a determined order. Penguins, however, renew all their feathers at the same time, within two to six weeks.



## SEASONAL CHANGE


In the high mountains, snow transforms the landscape during winter. During this time, nonmigratory birds exchange their summer plumage for a winter one. This change helps them to protect themselves from predators.

## PTARMIGAN



**SUMMER PLUMAGE**  
The feathers have deep pigmentation. This helps birds blend in with the vegetation.

**WINTER PLUMAGE**  
The new, unpigmented feathers make it possible for ptarmigans to blend with the white snow.



## Order of Replacement

Many species start molting, a process triggered by hormones, in a specific order. Molting starts with remiges and wing coverts, continues with rectrices, and finishes with body coverts. This gradual process keeps the body temperature stable.

- 4 Massive replacement of chest, back, and head coverts occurs from the center outward. This change coincides with the substitution of the seventh remex (singular of remiges).
- 3 Rectrices are replaced from the center outward. This happens simultaneously with the loss of tertiary remiges.

2 The wing coverts are replaced.

1 Renewal starts in the first primary remiges and spreads outward. In the secondary remiges, it spreads in two directions. Replacement occurs when the new remiges are three fourths developed.

61%

THE PERCENTAGE OF A BIRD'S BODY COVERED BY FEATHERS WHEN RENEWAL IS AT ITS PEAK

1 In the epidermal papilla, the formation of the new feather causes the detachment of the worn-out one.

2 A papilla develops from skin cells. The epidermal cells multiply faster than the dermal ones and form a collar-shaped depression called the follicle.

3 The papilla grows and becomes layered. The outermost layer is covered with keratin, which protects the underlying Malpighian layer (nucleus of the papilla). A group of dermal cells brings nutrients through the blood vessels that travel along the new feather.

4 The rapid growth of the Malpighian layer starts to develop the new feather. The rachis, barbs, and barbules become keratinized. The vessels that bring nutrients are reabsorbed, and the connection with the dermic layer is closed. Finally the protective vane breaks, and the feather unfurls.

5 The feather, now lifeless, assumes its characteristic blade shape. A residue of dermal and epidermal cells at the base of the follicle forms an area that will allow for replacement when the feather wears out.

**20 days**  
IS THE AVERAGE AMOUNT OF TIME THAT IT TAKES FOR A NEW FEATHER TO FORM.



# Gliding


Involves using air currents to fly and save energy when traveling long distances. There are two types of gliders, terrestrial birds and marine birds, each of which is adapted to different atmospheric phenomena. Terrestrial birds rise on thermals (rising air currents). Marine birds make use of oceanic surface winds. Once the birds gain altitude, they glide off in straight paths. They slowly lose altitude until encountering another thermal that will lift them. Both terrestrial and marine gliders have wings of considerable size. ●

## TYPES OF GLIDING FEATHERS

**Terrestrial Glider**  
A large wing surface allows it to make the most of rising air currents at moderate speed.



**Marine Glider**  
Thin and long wings allow it to make the most of the constant surface winds and offer less resistance to forward movement.



## Takeoff

Usually, a powerful jump followed by the vertical flapping of the wings is enough to make a bird take flight. As it descends, the tip feathers are stacked on top of each other, forming an airtight surface that helps drive the bird upward. As the bird raises its wings to repeat the movement, the feathers curve and open until the wing reaches its highest point. With a couple of flaps of the wings, the bird is in flight. Bigger birds need a running start on the ground or water in order to take off.



**70%**  
THE ENERGY SAVED BY A SEAGULL WHILE GLIDING

### SECONDARY FEATHERS

There are many of these because of the wing's length.

The wing length of some pelicans may reach **8 feet (240 cm)** from tip to tip.


### PRIMARY FEATHERS

There are fewer of these, as they only form the tip.

### WINGLETS


Terrestrial gliders usually have separate primary feathers (toward the tip of the wing) that serve to decrease the noise and tension generated there by the passing of air. Modern airplanes copy their design.

The tip feathers work as airplane winglets.



## Marine Birds

Dynamic soaring is performed by birds with long and thin wings, such as the albatross. These wings are designed to take advantage of horizontal air currents, which are responsible for the formation of waves in the ocean. The result is a flight consisting of a series of loops as the bird is lifted upward when it faces the wind and moved forward when it faces away from the wind. This kind of flight can be performed at any time.



### THE WING

Its particular shape causes lift, with its convex side and less pronounced concave side.

**FASTER AIRSTREAM**  
**CONSTANT AIRSTREAM**

LIFT

### PATAGIUM

Elastic and resistant skin covering with feathers. It is the wing's cutting edge, responsible for dividing the airstream.

### UPPER SIDE

Convex. The air covers more distance and accelerates, causing a lower pressure that "sucks" the wing upward.


### LOWER SIDE

Concave. The air covers less distance, it does not accelerate, and its pressure does not change.

**SPEED OF DISPLACEMENT** depends on the strength of the headwind.

## TERRESTRIAL BIRDS



They use warm, rising air currents generated through convection in the atmosphere or through the deflection of air currents against crags or mountains. Then they glide in a straight flight path. This type of flight is possible only during the day.



## FLIGHT PATTERNS

Flying in formation is a way for birds in flapping flight to save energy. The leader encounters more resistance as it flies, while the others take advantage of its


wake. There are two basic patterns: "L" and "V." The first is used by pelicans, and the second is used by geese.





# Flapping Flight

Most flying birds use flapping flight all the time. It consists of moving through the air as if rowing with the wings. With each flap (raising and lowering), the wing both sustains the bird in the air and pushes its body forward. There are different types of flapping flight and different rates of flapping. In general, the larger the bird, the more powerful and less frequent its flapping will be. Because flapping is an activity that consumes much energy, birds have adapted a variety of flight patterns: some, like hummingbirds, always flap their wings, whereas others alternate flapping with short-term gliding. The wing shape also varies according to the bird's needs. Birds that cover long distances have long, narrow wings; those that fly among trees have short, rounded wings. ●



**1**  
**Upstroke**  
As the wings move upward, the remiges separate and form grooves to reduce friction. Support for the bird comes from the patagium, a layer of skin that anchors the feathers and covers the bones.

**STRENGTH**  
To gain height above the ground, the wings flap in big arches in a manner that generally produces a lot of noise.

**2**  
**Downstroke**  
As the wings move downward, the remiges are forced together, and the wing moves forward a little for extra support. The wing also bends at the tips to push the bird forward, as if it were rowing.

Muscular strength is distributed to the entire wing, but it increases near the tip.

The downstroke of the wing provides propulsion.

**WING STROKE**  
The wing acts like an oar as it traps air and pushes the bird forward.

**THE CROP**  
Made of elastic skin. It can hold food during flight.

**THE LEGS**  
remain at rest until landing. They stay very close to the body.

**THE TAIL**  
Slightly curved, it works as a rudder during flight and as a brake during landing.

**THE BILL**  
Projected forward, its aerodynamic shape decreases the bird's air resistance.

**THE HEAD**  
Tilted backward to bring it closer to the center of gravity (between the wings) and attain balance

**ANGLE OF THE WING**  
Variable, depending on the wing's position. It closes on the downstroke.


**30 miles per hour (50 km/h)**  
**THE AVERAGE SPEED OF AN ADULT PELICAN DURING FLIGHT ON A WINDLESS DAY**

## A Specialized Design

Flapping flight is an activity that requires much effort. Therefore, birds must eat large amounts of food. A migrating swallow uses 4 kilocalories (4,000 calories) per 1.6 miles (2.5 km) of flight, whereas a small mammal needs only about 0.025 kilocalorie (25 calories) to travel the same distance.

## WINDMILL FLIGHT: HUMMINGBIRDS

Hummingbirds are able to hover in order to suck the nectar out of flowers. In contrast to other birds, hummingbirds' wings are attached only at the shoulders, which provides greater freedom of wing movement, allowing the hummingbird to hold itself in the air during both the upstroke and the downstroke. The hummingbird has to flap its wings up to 4,800 times per minute during directional flight and for hovering.



The wings flap 80 times per second during normal flying.

**Courtship Display**  
Certain hummingbird species can flap their wings up to 200 times per second during courtship.



The wing has short, sturdy bones; the muscles are very powerful.

**Great Maneuverability:**  
Hummingbirds are the only birds capable of moving backward.

## Landing

requires reducing speed until the bird becomes motionless and settles. The bird faces the wind and spreads out its tail, wings, and alulae (bastard wings, characterized by their stiffness and growth from the first digit), while lifting up its body and extending its legs forward to increase the surface area in contact with the air. In addition, the bird flaps its wings intensely in the direction opposite to its flight. Everything works like an aerodynamic brake. Some birds—such as the albatross, with its long, narrow wings—tend to have problems slowing down. As a result, they are ungainly when landing on the ground, but on the water they are able to ski on their feet until coming to a stop.



Flapping Against the Wind

Open Wings

Spread Tail


Sliding

The feet spread open before landing to provide more resistance and help the bird to slow down.

## WAVELIKE FLIGHT PATH

Ideal for high speeds, it consists of flapping the wings to gain height and then folding them in order to descend along the flight's trajectory. Afterward the bird flaps its

wings again, making use of the inertia of its descent to regain height. A variation of this type of flight involves gliding between flaps of the wings.



**1** Propulsion  
The bird flaps its wings to ascend.

**2** Rest  
The bird keeps the wings near its body to save energy for short intervals.

Ascent

Flapping Wings

Folded-up Wings

Descent



# Speed Records

The world of birds is amazing when expressed in numbers. Most birds travel at speeds between 25 and 45 miles per hour (40–70 km/h), but when diving, peregrine falcons can reach more than 200 miles per hour (320 km/h). Many species can reach an altitude of 6,600 feet (2,000 m), although climbers have seen geese flying over the Himalayas at more than 26,000 feet (8,000 m). The fastest swimmer is the Gentoo penguin, which can swim 22.4 miles per hour (36 km/h). Considering its small size, it is surprising that the *Selasphorus rufus*, or Rufous hummingbird, which is only 4 inches (10 cm) long, carries out an extensive round-trip migration each year from northern Alaska to Mexico. Here are some more incredible facts. ●

**RUPPELL'S GRIFFON VULTURE**  
**36,870 feet**  
 (11,237 m)

In 1973 a Ruppell's griffon vulture crashed into an airplane flying over the Ivory Coast at this altitude.



Scale (in thousands of feet)

Weight 15-20 pounds (7-9 kg)

Flying Altitude commonly reaches 20,000 feet (6,000 m).

## Air

### SPEED ▶

Most birds fly between 25 and 45 miles per hour (40 and 70 km/h), but the fastest birds can beat the cheetah, the most famous of the fast animals.

### DRAGONFLY


The fastest flying insect. It reaches

**31 mph**  
 (50 km/h).



### ALTITUDE ▶

Flying at high altitudes requires a strengthened circulatory system to make up for the scarcity of oxygen in the air.



**BLACK SWAN**  
**27,000 FEET (8,230 M) OF ALTITUDE,**  
 according to a pilot who witnessed a flock over the Hebrides Islands


**BAR-HEADED GOOSE**  
**28,000 FEET (8,500 M)**  
 Some climbers reported having seen specimens of geese flying at 28,000 feet (8,500 m) over the Himalayas.

**CHOUGH**  
**29,030 FEET (8,848 M)**  
 A group of climbers on Mt. Everest found choughs standing on the summit.

## Land


**ELEPHANT**  
 17 MPH (28 KM/H)

Scale (in miles per hour) Land-Water



**PEREGRINE FALCON**  
**200 mph**  
 (320 km/h)  
 FASTEST BIRD WHILE DIVING

Weight of Males 1.1-2.4 pounds (0.5-1.1 kg)  
 Weight of Females 1.5-3.5 pounds (0.7-1.6 kg)



## Water

**DOLPHIN**  
 22 MPH (35 KM/H)

**SEI WHALE**  
 30 MPH (48 KM/H)  
 Fastest swimming mammal

**TUNA**  
**62 mph**  
 (100 km/h)  
 FASTEST SWIMMING FISH OVER SHORT DISTANCES

**CHEETAH**  
**65 mph**  
 (105 km/h)  
 FASTEST MAMMAL OVER SHORT DISTANCES

### DISTANCE I

The arctic tern travels **24,850 miles** (40,000 km).

It migrates from Canada and Labrador to Antarctica and the Austral Sea. On each trip, it travels 9,000 to 12,000 miles (15,000-20,000 km).

**ARCTIC TERN**  
 (*Sterna paradisaea*)




### DISTANCE II

The Rufous hummingbird flies from northern Alaska to Mexico and back—a journey of

**6,000 miles**  
 (10,000 km/h).

**RUFOUS HUMMINGBIRD**  
 (*Selasphorus rufus*)

Weight 0.1-0.2 ounce (4-6 g)



### ENDURANCE

The endurance record goes to the golden plover, which is able to fly a distance of **1,900 miles** (3,000 km) without stopping.

**GOLDEN PLOVER**  
 (*Pluvialis apricaria*)





# The Lives of Birds

**PARTRIDGE EGGS** (*Lagopus lagopus scoticus*)  
The female lays eggs at intervals of one to two days, and she is the one who incubates them.

**THE ANNUAL CYCLE** 42-43

**HOW THEY COMMUNICATE** 44-45

**NUPTIAL PARADE** 46-47

**HOME SWEET HOME** 48-49

**FIRST, THE EGG** 50-51

**BIRTH IN DETAIL** 52-53

**POSTNATAL DEVELOPMENT** 54-55

**A DIET FOR FLYING** 56-57

**MIGRATION ROUTES** 58-59

**DEFENSE STRATEGIES** 60-61



**T**he behavior of birds is closely connected to the seasons. To survive, birds must prepare for the arrival of fall and winter and adjust their behavior

accordingly. Gliding over the oceans, a wandering albatross, for example, can travel anywhere from 1,800 to 9,300 miles (2,900 to 15,000 km) in a single day in search of food. When the time comes

to choose a partner, the behavior of males is different from that of females: males employ a variety of tactics to win over females and convince them of their fitness. Some bird couples stay together

forever, whereas other birds change partners every year. As for caring for chicks and building nests, in most species both parents participate. ●




# The Annual Cycle

The annual cycle of seasons is like the daily cycle of night and day. Fluctuations in the intensity of light over time create a series of physiological and behavioral changes in birds, whether throughout the year or throughout the day. This biological clock is clearly reflected in birds' reproduction and migrations. Changes in light that are detected by a bird's retinas induce the secretion of melatonin by the pineal gland. The blood level of this hormone acts on the hypothalamus-hypophysis axis, which regulates internal processes. This is one reason why birds start to change their plumage and feel the need to fly to other areas. ●

## How the Hypophysis Works

Reproduction is the main activity under the control of the hypophysis, which determines behaviors such as finding a place to court females and mate, building a nest, incubating the eggs, and stimulating unborn chicks to break their shells. The hypophysis is a gland in the brain that has several functions. It receives nervous and chemical

stimuli and produces hormones. These hormones regulate the metabolic activities that cause birds' internal and external sexual organs to develop. For example, the gonads become enlarged, and secondary sexual characteristics, such as ornamental crests or plumes, appear.



### Sleep Regulator

The pineal gland, or epiphysis, produces melatonin. The level of this hormone determines the phases of sleep and wakefulness.

### THE MOST IMPORTANT GLAND

The hypophysis is located in the ventral area of the brain, below the hypothalamus. Its secretions control vital functions, from blood pressure and the balance of water and salts in the body to the activity of the gonads and the thyroid.

### ANNUAL CYCLE

Incubation, migration, and courtship activities vary according to the amount of light available during each season.

1



In early summer, incubation takes place. The increase in the amount of daylight coincides with this phase of the annual cycle.

2



The arrival of the first days of fall coincides with a decrease in the amount of light. The migration season begins.

3



The amount of light increases with the beginning of spring; males use their huge throat pouches to court females.

## Survival Manual

Birds' most striking behaviors are associated with the reproductive season. During courtship parades, birds engage in elaborate choreographies; there are also extraordinary fights between males. The blue-footed booby, the male frigate bird, and the ruff are just a few examples of birds that engage in these behaviors. Others, such as the snowy egret (*Egretta thula*), prefer to offer twigs for the construction of the nest. The Vogelkop bowerbird (*Amblyornis inornata*) builds bowers with leaves, flowers, or any other object that may help him to win over the female. Birds' performances are not connected only to courtship. The killdeer (*Charadrius vociferus*) fakes being wounded to defend the eggs or chicks in the nest from predators. It offers itself as easy prey by dragging a wing as if it were broken. This trick shifts the danger away from the young.

### SHOWING OFF

The magnificent frigate bird (*Fregata magnificens*) is a large bird that lives in coastal areas. It has large wings, powerful talons, and a strong hooked

bill. During the reproductive season, it is responsible for building the nest. With its impressive appearance, it endeavors to attract a female.



#### New House

While it prepares the nest, its chest rests, with its skin pink and relaxed.



#### Red Chest


The throat pouch remains inflated for several hours or until the female chooses the most seductive male.

#### Indicating Repose

To rest, the pelican reclines its head and places its bill under a wing.

### DANCE OF THE BLUE-FOOTED BOOBY

(*Sula nebouxi*) The males—and, on occasion, the females—perform a graceful courtship dance after marking the territory for nesting. They sing and show off their plumage with careful synchronization.



1 **Raised Head**  
It flaps its wings and marches, looking at the sky.

2 **Parade**  
It lowers its head and parades like a soldier around the nest. Finally, it shakes its whole body.

### IN COMBAT

In the summer, male ruffs develop a huge "ruff" and auricular feathers around their necks. Their courtships are violent and striking. When competing for mating territory, they struggle fiercely. Afterward they docilely sprawl their bodies on the ground until the female chooses the lucky one.



Male Ruff  
*Philomachus pugnax*

TO THE ORGANS




# How They Communicate

Sound is an important form of expression in the lives of birds. Birds' sounds can be of two types: calls and songs. The former have a simple acoustic structure, with few notes. They are associated with coordinating the activities of a group, establishing communication between parents and their young, and maintaining contact between birds during migration. Songs, on the other hand, are more complex in rhythm and modulation. They are controlled by the sex hormones, primarily the male hormones. For this reason, males produce the most varied melodies. Songs are linked to sexual behavior and territorial defense. In general, birds either inherit or learn them. ●

## Territoriality and Range


One of the most studied functions of birds' songs is territorial demarcation. When a bird occupies a territory, it sings to announce its claim to competitors, as the pipit shown to the left is doing. When birds must share territory, as in a colony, they develop dialects (variations of sounds produced by the species). When a bird

born and raised in one location moves, it must learn the dialect of the new location in order to be accepted and participate in the community. There are also mechanical sounds produced by wing strokes, legs, and bills. In a display of territorial defense, the eared nightjar combines singing with beating its wings.



4,000 bird species


SHARE WITH HUMAN BEINGS AND WHALES THE NEED FOR "SOMEBODY" TO TEACH THEM TO VOCALIZE (SONGBIRDS, HUMMINGBIRDS, AND PARROTS ARE EXAMPLES).



**INTENSITY** can vary widely from bird to bird. The larger the territory, the greater its reach. Its frequencies can change as well: the lower the frequency, the greater the coverage.


### 1 THE SONG AND THE BRAIN

Birds have a brain that is well developed for this function. Testosterone acts on the upper vocal center of the brain, which is in charge of memorizing, identifying, and transmitting the orders for the execution of the song.



### 2 EXPULSION OF AIR TO THE BRONCHI

The air stored in the air sacs and lungs is expelled. As it passes through the syrinx (located between the bronchi and the trachea), it vibrates the tympaniform membranes. These membranes are the equivalent of vocal cords in humans.




### 3 THE PRODUCTION OF SOUND IN THE SYRINX

The participation of both the sternotracheal muscles and five to seven pairs of small internal muscles is needed for producing sounds. These muscles control the elongation and contraction of the syrinx, which varies the pitch of the sound. The air sac is also important because it adds external pressure, which causes the tympaniform membranes to tighten. The esophagus works like a resonating box, amplifying the sound. The articulation of the sounds occurs in the buccopharyngeal cavity. There are two types of articulation: guttural and lingual.

#### SIMPLE SYRINX


The tympaniform membranes are located above the place where the bronchi divide. They are moved by a pair of external muscles.



#### SONG-PRODUCING SYRINX


##### A AIR AND BRONCHI

During breathing, the songbird maintains airflow without affecting the resting syrinx.




##### B CLOSED MEMBRANE

The membranes close on both sides, under pressure from the external muscles. The bronchi rise slightly and also adjust the membranes.



##### C THE SOUND

The membranes vibrate with the air current and propagate the sound through the trachea until it reaches the bird's bill.



## Strengthening Ties


Some songbirds develop very complex singing rituals. The duet is perhaps the most striking because it requires both a shared repertoire and good coordination between both birds. In general, the male initiates the song with a repeated introduction; the female then alternates with different phrasing. The phrasing allows for more or less cyclical variations that make it unique. It is believed that this strengthens ties between the pair (as demarcation of territory does) and serves as a stimulus for cooperative behaviors, such as nest building, in which both the male and female may participate.





# Nuptial Parade

Finding a mate is not easy for any species. For birds, the exhibition of plumage with bright colors, the presentation of offerings and gifts, and the performance of dances and highly elaborate flight patterns are some of the particular behaviors seen during this period. They are known as nuptial or courtship displays. The male resorts to all these strategic gestures to attract the female's attention and prevent her from paying attention to other males. Some of these rituals are extremely complicated; others are very tender and delicate. ●



## AERIAL EXHIBITIONS

Certain birds, such as goshawks or male northern harriers, court the female in flight. They ascend in the air in broad circles, only to let themselves fall in daring, sharp dives.



## DISPLAYING PHYSICAL ATTRIBUTES

To find a partner, birds such as the snowy egret resort to a series of very elaborate signals, such as songs, poses, dances, flight patterns, noisemaking, and displays of their ornamental feathers.



## BUILDING BOWERS

Australian bowerbirds build a structure called a bower, which they decorate with pieces of paper and fabric that inevitably attract the female.



## GIFTS

Another courtship strategy is the presentation of gifts. Male eagles give females prey, and European bee-eaters offer insects. These offerings are called courtship food.

## MUTUAL DANCES AND COURTSIPS

Great crested grebes (*Podiceps cristatus*) perform incredible aquatic dances. They bow to each other, dive, and run through the water side by side.

## Special Courtship

Avian courtship is a phenomenon that, depending on the species, can take the form of various rituals. Lek rituals are one of the most intriguing forms of courtship. The males gather in a small area, called an arena, where they perform their courtship displays for the females. The females form a circle around the arena and end up mating with the male that has the most striking secondary sexual characteristics. Lek is a system controlled by the dominant male, who ends up mating with most of the females (polygyny). The less experienced males will mate with only a few, or even none, of the females. For some species, lek rituals can be very intricate. At least 85 species perform this special type of courtship ritual, among them manakins, pheasants, cotingas, and hummingbirds. Manakins, for example, stand in line and wait their turn to perform.



## TIMING

The courtship display is directly related to reproduction cycles. It takes place before copulation, although it can continue to occur thereafter.

**Prenuptial**  
Prenuptial courtship starts with territorial establishment and the search for a partner, which can take place simultaneously.



**Postnuptial**  
With this display, the great crested grebe ensures the continuity of the pairing even after the eggs are laid.



5.9 feet  
(1.8 m)


IS THE SIZE OF THE TAIL OF THE PEACOCK WHEN IT UNFURLS ITS MORE THAN 200 SHINING FEATHERS AND FORMS A FAN TO ATTRACT THE FEMALE.

**Gray Crowned Crane**  
*Balearica regulorum*  
Two cranes perform a courtship dance consisting of a series of impressive leaps.

**Emperor Penguin**  
*Aptenodytes forsteri*  
A monogamous species. Each recognizes its partner by its voice, and couples will spend their entire lives together.

## Monogamy or Polygamy

Monogamy is the most common mating system, in which two birds, one of each sex, participate, leading to the formation of a couple. This couple can endure for a single reproductive season or for life. Polygamy is an alternative pattern, but it is not very common. Polygamy is divided into two classes: polygyny, in which the male mates with several females, and polyandry, in which the female mates with several males (and may even be able to keep them all together in a harem). In either case, one partner has the sole responsibility of caring for the eggs and chicks. There is also an exceptional case within polygamy: promiscuity. In this arrangement, a couple is not formed, and the relationship is limited to copulation.






# Home Sweet Home

Most birds lay their eggs in a nest, where they are incubated by the body heat of an adult sitting on them. To build a nest, the couple normally uses mud mixed with saliva, small stones, branches, and feathers. When the nest is in a visible location, the bird covers it with lichens or loose twigs to hide it from predators. Nest shapes vary according to the bird group: they can take the form of a bowl, a hole in a tree (woodpeckers), or an excavated burrow on a slope of sand or soil. There are even birds that use nests built by other species. ●

## SOME VARIATIONS




**WOVEN NESTS**  
Weaver birds intertwine grass blades until they form a structure. The entrances are underneath.



**BURROWS**  
Parrots and kingfishers dig their nests in sandy river banks.




**SEWN NESTS**  
The tailorbird sews two large leaves together with grass blades. The nest is inside the leaves.



**PLATFORM NESTS**  
The sparrow hawk gathers a large number of branches and assembles a high, solid base for its eggs.

## Types and Locations


Nests are classified according to their shape, material, and location. They vary depending on the amount of warmth the species needs, as well as in terms of the level of protection they offer. The greater the pressure from predators, the higher or better hidden a nest must be. Good examples are isolated nests resembling high platforms; nests in deep depressions in the soil or hidden in tree trunks, which are very safe and provide good insulation; and nests made of clay, which are very hard. The most typical nests resemble a cup and are found at various locations, most often between two or three high, remote branches.




## How the Nest Is Built

A cup-shaped nest is built at a fork between two or three branches. The bird arranges twigs, grass blades, and small sticks, as if building a platform. The bird then interlaces some of these materials with the tree to give the nest solidity. It then interweaves the materials in a circular pattern. As the nest takes shape, lighter, more adhesive materials—such as mud, spiderwebs, caterpillar silk, and certain plant fibers—are used. Although the outside is rough, the inside is lined with feathers for softness and warmth. In general, if both the male and female participate in the construction, a few hundred trips are enough to complete the nest. In some species, such as weavers, males have to display a nest during courtship; in others (African black eagles, for instance), the same nest is used every year.


**THE LITTLE MASKED WEAVER** builds solid nests, weaving together leaves and grass blades. Sometimes the male builds several nests before the female chooses him.



**1 BASE**  
The bird gathers branches and small sticks on a fork to lay the foundations of the nest, setting up a platform-like structure. The bird then ties it to the tree, making it sturdy.



**2 SHAPE**  
The bird intertwines grass blades, small sticks, and hairs; fastens them; and gives them a circular shape. As construction advances, the bird uses adhesive materials, such as spiderwebs.



**3 END**  
The finish includes moss and feathers to smooth out the interior and provide insulation against the wind and cold, which makes the nest more suitable for incubation.

**BASE**  
This is the nest's foundation, and it is the first section built by the bird. It is solid and is usually built with bulkier materials.

**WALLS**  
Are the most important component in a nest's construction, and they give it its characteristic shape. The materials used vary, depending on the habitat.

**LINING**  
It is composed of fibers, hairs, feathers, and down. It insulates the eggs from the cold and helps with incubation.

## Structure

The cup shape is important for preventing eggs from rolling out and falling. Besides making the construction of the nest easier, the use of different materials helps make the nest sturdier, as the smaller and more flexible pieces firm up the base, walls, and lining of the structure. Different materials also provide more efficient insulation, keeping warmth in and cold out during the incubation and raising of chicks. As an added reinforcement, birds usually make the side facing the prevailing winds thicker and the side facing the sun thinner. Thus, the whole nest is an incubator that conserves energy. Finally, an exterior finish camouflages the nest among branches, hiding it from predators.






# First, the Egg



Birds may have inherited their reproductive method from their predecessors, the theropod reptiles. In general, they lay as many eggs as they can care for until the chicks become independent. Highly adapted to the environment, the eggs of the same species have varying shapes and colors. These variations help keep them safe from predators. They also vary greatly in size: the egg of an ostrich is 2,000 times bigger than that of a hummingbird. ●

## How It Forms


Birds have only one functional ovary, the left one, which grows dramatically during the mating season. The ovule can descend and form what are known as unfertilized eggs (the type used in cooking). If the egg is fertilized, embryonic development begins. The ovule, fertilized or not, descends to the cloaca in a few hours or days. The eggshell begins to be formed at the isthmus, through the secretion of calcium. At first soft, the shell hardens when it comes in contact with the air.




**LAYING**  
A group of eggs laid at one time is called a laying. During the mating season, a sparrow can have several layings. If some eggs are removed, the sparrow can replace them without difficulty.




**SHAPE**  
It depends on the pressure exerted by the oviduct walls. The large end emerges first.



**SIZE**  
There is no exact proportion between the size of a bird and its egg.



**THE SHELL**  
Formed by a solid layer of calcium carbonate (calcite), it has pores that make it possible for the chick to breathe. Bacteria are kept out by two membranes that cover the egg, one on the inside and the other on the outside.



**8%**  
THE PROPORTION OF AN EGG TAKEN UP BY THE EGG SHELL

**ALBUMIN**  
was consumed.  
**YOLK**  
disappears into the body.




# Birth in Detail

When a chick is about to hatch, it starts to make itself heard from inside the egg. This allows it to communicate with its parents. It then starts to peck at the shell with its tiny egg tooth, which is lost after birth. Next, it turns inside the egg and opens a crack with new perforations, at the same time pushing with its neck and legs until it manages to stick out its head. This job demands a lot of effort and can take 30 to 40 minutes or, in the case of kiwis and albatrosses, even three to four days. In most species, newborn chicks are blind and naked, and they can open their bills only to receive food. ●

## INCUBATION

For the embryo to develop, it needs constant temperatures between 99° and 100° F (37-38° C). The parents ensure these temperatures by sitting on the eggs and warming them with their brood patches.



During incubation, some species lose their chest feathers and increase their number of blood vessels in this area. Others pluck out their feathers. Direct contact with the eggs helps keep them warm.

## DURATION BY SPECIES

The incubation period varies considerably: between 10 and 80 days, depending on the species.



**PIGEON**  
Females and males incubate. They both develop a brood patch.  
**18 days**



**PENGUIN**  
Both males and females incubate. The emperor male has a special pouch for incubation.  
**62 days**



**ALBATROSS**  
Lacking brood patches, the parents hold the egg between their feet and abdomen.  
**80 days**

## Breaking the Shell

This process may take from a few minutes to three or four days, depending on the species. In general, the parents do not intervene or help their young. When the shell is empty, they throw it out of the nest, apparently to avoid attracting the attention of predators. In species whose young

hatch with the feathers already developed, hatching is extremely important. It has been observed that the singing of the chicks stimulates the stragglers and delays those that have gotten ahead; it is important that they all be ready to leave the nest together.

**35 minutes**

IS THE APPROXIMATE TIME IT TAKES A SPARROW TO COME OUT OF THE EGG.

**1**

### CRACK IN THE EGG

The chick turns inside until its bill targets the egg's midline. It then punctures the air sac. With a few more tries, it pierces the shell. The chick then breathes for the first time.

### ASKING FOR HELP

The chick calls for its parents from inside. The reply encourages it to continue the effort.



### SEQUENCE OF PECKING

Between each sequence of pecking, the chick must take long breaks.



**2**

### THE CRACK EXPANDS

After making a hole in the shell, the chick opens a crack with successive pecks at other points. Air gets in and dries up the membrane, which makes the task easier.

## Adaptations for Hatching

Getting out of the egg is an intricate operation because the space is tight, and a chick's muscles have little vigor. Birds count on a few adaptations, such as the egg tooth and the hatching muscle, to accomplish the task. The tooth is used for making the first perforation, which allows air into the egg. The muscle exerts the necessary strength, while stimulating the chicken's motor functions to intensify the effort. Both the egg tooth and the hatching muscle disappear shortly after the eggshell is broken.

### HATCHING MUSCLE

It exerts pressure against the shell and helps to break it.



### EGG TOOTH

A protuberance on the bill that punctures the egg. Its presence depends on the species.




**4**

### THE CHICK IS BORN

Once outside the chick, almost featherless, looks for warmth and food from its parents. In the case of some birds that hatch without feathers, not all eggs hatch simultaneously; this benefits the firstborn if food is scarce.

### A GREAT EFFORT

Getting out of the shell requires much energy from the chick.



SHELL MEMBRANE

EGGSHELL


**3**

### GETTING OUT OF THE EGG

Once the shell is open, the chick pushes itself out with its legs and by crawling on its abdomen. For birds that hatch without feathers, this is more difficult, because they are less developed.

### WHAT COMES OUT FIRST?

The head usually comes out first, because the sharp bill helps break the shell. Most birds then get out of the egg by pushing themselves out with their legs. For wading birds and other terrestrial birds, however, the wings usually unfold first.





# Postnatal Development

Chicks develop at highly variable rates after hatching. Some birds are born with their eyes open and with a thick layer of down feathers. These birds can also feed themselves. That is why they are called precocious, or nidifugous. Ducks, rheas, ostriches, and certain beach birds can walk and swim as soon as they are born. Other species are born naked and develop their feathers later. They need to stay in the nest until they have sufficiently developed, so adult birds must care for them. These birds are called nidicolous. The most helpless chicks are the young of songbirds and hummingbirds, because they need warmth from their parents to grow strong. ●

## Nidifugous Young

Nidifugous young are fully developed at the moment of birth. They can move and even leave the nest, hence their name (which means "fleeing the nest"). This adaptation demands more incubation time because the chick is almost fully developed at birth. This is the case with incubator birds (*Megapodius freycinet*), which begin their independent lives in the outside world as soon as they leave the shell. Ducks follow their parents but find food on their own, whereas chickens follow their parents, and the parents show them where to find food.

### GROWTH STAGES


**24 hours**

IS THE MINIMUM AMOUNT OF TIME THE BLACK-HEADED DUCK NEEDS TO BE READY TO FLY.



**7 to 8 days**

Growth quickens, and the first covert feathers appear at the tip of the wing. The bird leaves the nest. Its diet consists of: **66% invertebrates.** The rest consists of seeds and flowers.



**Red-Legged Partridge**  
*Alectoris rufa*

**MOVEMENT**  
Within a few hours of hatching, nidifugous birds can run around.

**FEATHERS**  
The body emerges from the egg covered with damp down. Within three hours, it will become dry and fluffy.



**EYE**  
They are born with open eyes.

**21 days**

It is already considered an adult. Its flights are longer. Its diet is composed of 97 percent vegetables; the rest is made up of lichens and insects.

### SIZE COMPARISON

#### NIDIFUGOUS

The egg is larger, the chick is born more developed, and the incubation period is longer than that of nidicolous birds.



#### NIDICOLOUS

They lay small eggs, with a brief incubation period, and the young are helpless at birth.



## Nidicolous Broods

Most of these chicks are born naked, with closed eyes and with only enough strength to get out of their shells. They stay in the nest. For the first few days, they cannot even regulate their own body temperature; they need their parents in order to stay warm. Within one week, they have a few feathers, but they require constant care and food. They form a numerically important group that includes Passeriformes (songbirds).

**FOOD**  
They need much food to develop. The parents must feed them 24 hours a day.




**House Sparrow**  
*Passer domesticus*

An adult bird can feed its young up to **400 times a day.**

### INTERIOR OF THE BILL

Its color is bright to stimulate the parents to regurgitate the food.





**Shining Areas**  
Some species have shiny areas that can be seen even in the dark.

### EYES

Nidicolous chicks are born blind. They open their eyes a few days after birth.

### FEATHERS

The chicks are born either naked or with down feathers in some areas.



**12-15 days**

Development is complete, and the covert feathers are ready for flight. All that remains is for the bird to reach adult size.



**10 days**

Feathers cover everything, but they are not yet developed. The chick can stay warm on its own, and it is voracious. Growth is very fast.



**8 days**

Feathers cover the chick almost completely, except around its eyes. Its legs are well developed, and the sparrow moves around in the nest.

### STAGES OF GROWTH


**25 hours**

It performs a few instinctive movements. It can barely raise its head to ask for food.



**4 days**

The eyes open. The tips of the first feathers appear. It performs a few movements.



**6 days**

Some feathers begin to unfold, the nails are formed, and the wings continue to grow with the body. It can stand up.

**12-15 days**

IS THE ESTIMATED TIME THAT IT TAKES THIS NIDICOLOUS CHICK TO LEAVE THE NEST.




# A Diet for Flying

Most birds eat assorted foods that are rich in energy and proteins. Their high level of activity requires that they eat almost constantly. Their sources of food are varied and include seeds, fruits, nectar, leaves, insects and other invertebrates, and meat of all kinds (including carrion). Many species eat more than one type of food; some even alternate according to the seasons and to the migration cycle. This guarantees their survival. However, there are others birds—a minority—that consume only one type of food, for which they have no competition. Because their dependence on this single source of food is greater, though, the risk is higher. Feeding behavior also varies among different species. For example, some eat alone, and others eat in groups. ●

## A Complex System

Feeding on microorganisms that live in salty water demands a complex filtration system. The flamingo's bill is specially suited to this task. Its tongue and throat pump the water inside the bill as they ascend and descend, bringing water through the hornlike lamellae, which resemble whale baleen, to retain the food passing through them. Their tongues also have a depressed area, where the stones and sediments that come in with the water collect. The whole operation requires that the flamingo submerge its bill upside down. Because of the number of microorganisms they need to consume and the time it takes to filter them, flamingos usually spend many hours in the water. The risks involved in this activity are mitigated by the fact that flamingos do not eat alone, but rather in groups. Occasionally, there are instances of aggression, possibly because of territorial conflicts.

### Filtration of Food



## From Parents to Children

Flamingos and pigeons feed their young a special "milk" that is produced in the crop and has a nutritional value similar to the milk of mammals. Both males and females produce it as soon as food is ingested so that the chick is not fed regurgitated food.

It is given to the young through the bill. Easy to digest and nutritious, it is an excellent food.


The flamingo produces milk in its crop only when it is not eating. A protein called prolactin participates in its preparation, just as with mammals.

This milk has high concentrations of the pigment that colors the feathers; that is why the bird's color changes to the characteristic pink while molting in adolescence.

## Types of Diet

Birds expend a great deal of energy, and their diet must be suited to their high metabolic rates. They eat practically anything, although the diet of some birds is very specific. The same foods are not always available, which causes most birds to adjust their diets throughout the year.

**THE HUMMINGBIRD'S TONGUE**  
Long channeled or tubular tongues suck the nectar and catch insects at the bottom of the corolla of a flower.



**NECTAR**  
is a solution of sugar and water that flowers produce. It is very high in energy and easy to digest. In order to get it, a bird must have a long, sharp bill. In temperate regions, nectar is plentiful in the spring and summer, whereas in the tropical regions, it is available year round. Hummingbirds and honeycreepers are very fond of this juice.



**SEEDS AND GRAINS**  
The high energy and protein content of seeds make them an ideal food for birds. The problem is that they are seasonal.



**MEAT**  
Hunting and scavenging birds feed on meat. They rarely specialize in one type of prey, but the prey's size determines its selection.



**FISH**  
The abundance of fish throughout the year makes them one of the favorite foods of marine birds. They have a high nutritional value.



**LEAVES AND PLANTS**  
Few birds feed on leaves because they have low energy value. Birds that do have undergone adaptations that permit them to digest cellulose.



**FRUIT**  
In the tropics, this diet is very common because fruits are available throughout the year. In temperate and cold regions, fruit can be found only in the summer. Fruit has a high energy value, and many birds eat it.



**INSECTIVORES**  
Insects are rich in energy and proteins, and they are highly abundant. Thus, many bird species eat them. In cold regions, they can be found only in the summer.


## Strategies

Depending on the abundance of resources, the needs of the species, and the strategies for getting food, birds may eat alone or in groups. If food is scarce or widely scattered throughout a region, birds eat alone and defend their territory. On the other hand, if food is abundant, they prefer to eat within the safe confines of the group.




**SOLITARY**  
Usually birds of prey, such as eagles and owls, hunt alone because food is scarce and is randomly distributed. Hunting alone has a drawback in that birds must also watch out for predators, which takes time away from hunting.

**TOGETHER**  
This behavior is typical of ocean birds, such as pelicans and seagulls, and of aquatic birds, such as flamingos. When birds eat in groups, each group member can warn the others about possible dangers, which is beneficial.





# Migration Routes



**RUBY-THROATED HUMMINGBIRD**  
*Archilochus colubris*  
**500 miles (800 km)**  
 IS THE DISTANCE THEY FLY NONSTOP ACROSS THE GULF OF MEXICO. THEY DO IT IN JUST 20 HOURS.

**MEETING POINT**  
**1 billion**  
 BIRDS MEET IN THE DEAD SEA VALLEY EVERY YEAR WHERE THE ASIAN, EUROPEAN, AND AFRICAN MIGRATION ROUTES CONVERGE.

**ARCTIC TERN**  
*Sterna paradisaea*  
**25,000 miles (40,000 km)**  
 IS THE DISTANCE IT CAN COVER ON ITS ROUND-TRIP MIGRATION BETWEEN THE POLES. IT IS THE LONGEST MIGRATION IN THE WORLD.

## Behavior

In order to survive, millions of birds from all over the world start a journey every fall in search of better climates. The instinct of migrating, which other animals share with birds, was acquired after a long evolutionary process. Some birds fly thousands of miles through mountain ranges; others follow the course of a river without stopping until they reach their destination; others land after a short trip. In general, birds undergo physiological changes that are related to the length of their journey; some even lose almost half their weight during the migration. Routes tend to be fixed, although some migratory birds do not always follow them exactly. The colors on the map indicate the most important routes. There are collective and individual migrations, and birds may travel during the day or at night. It is surprising how fast birds travel. Many studies have recorded that carrier pigeons and white-headed sparrows can cover more than 600 miles (1,000 km) in a day. Shorebirds, such as godwits and curlews, are among the birds that cover the most distance. They are also among the birds most faithful to routes.


## Types of Migration

The north-south path traveled by many species is called latitudinal migration. A second pattern, which occurs in an east-west direction, is called longitudinal migration. The last pattern is called altitudinal migration; it is performed by birds that move up and down mountains seasonally.



## How They Find Their Way Around

Birds use a compass-and-triangulation system that lets them know where they are according to the position of the sun or stars. This system is similar to the one used by seafarers. It consists of measuring the sun's angle of inclination in relation to the horizon (azimuthal angle) and comparing it to the angle the bird knows through its biological clock. Birds also orient themselves by using the Earth's magnetic field. Furthermore, those migrating during the day register landmarks on their routes, such as mountains, lakes, or deserts. Still others follow older birds or are guided by their sense of smell.





# Defense Strategies

Birds have many predators, including cats, snakes, crocodiles, and other birds. To defend themselves against these predators, birds use various strategies, the most common of which is camouflage. Some birds blend in with their surroundings and thus go unnoticed by their enemies. Chaparral birds, whose plumage colors and patterns make them difficult to discern when they are on the ground, use this strategy. Other birds take flight in the face of a threat. There are also those that keep still in the presence of unknown animals, feigning death, whereas others face an enemy and fight. It is not uncommon to see magpies, thrushes, and other birds chasing away strangers that get close to their nests. ●

## Individual Strategies

Among solitary birds, it is common to flee quickly if a stranger is present. Not all solitary birds react this way, though; some have developed specific techniques to defend themselves.


### ESCAPE

In the presence of terrestrial predators, a bird's first reaction is to take flight. If the bird cannot fly, it looks for shelter or a hiding place.



### EXPANSION

Owls spread out their wings to look bigger than they really are.



### DISTRACTION

Little bustards shoot their excrement in the face of birds that prey on them. This distracts the predator and makes it possible to escape.



### CAMOUFLAGE

is very common and is one of the most efficient defensive strategies. Many birds develop plumage to imitate the dominant colors and shapes of the environment where they live. When they notice the presence of a potential enemy, they stay motionless to avoid calling attention to themselves. There are a few notable cases, such as that of the tawny frogmouth (*Podargus strigoides*), pictured above. Many partridges and terrestrial birds are experts at the art of blending in with the landscape; the rock ptarmigan, for example, has white plumage in the winter that becomes terra-cotta in the summer.

## Collective Strategies

Birds that have group behaviors usually develop group strategies to protect themselves against predators. Being numerous is a guarantee that the species will go on. They also adopt other tactics as a group.

### COLONIES

A great number of birds can defend themselves better from predators when in groups. For that reason they even form colonies with other bird species when raising their young.

### FLOCK

In the presence of predators, birds form flocks that fly in a synchronized manner, which makes it hard for the predators to focus on any one individual.

### ATTACKS AND COLLECTIVE AID

Many birds that live in groups have developed several hounding behaviors in the presence of potential enemies. They perform them to help an individual that is in danger or is unable to flee.

### WARNING

They emit callings that warn the whole colony. The great majority of species have a specific and characteristic cry that is usually simple, brief, and very audible. They often emit these warnings while adopting postures (such as stretching the neck or shaking the wings) that alone are enough to warn other individuals of the intruder's presence.

## Defense of the Brood

Hatching and youth are times of critical vulnerability for birds. During these phases, they are forced to keep still because they are easy prey. For this reason, parents permanently watch over their nests, even attacking strangers if they get too close.



### FEIGNING INJURY

A very widespread behavior is to pretend to be wounded or sick. This allows a bird to avoid being chosen as prey. This behavior is common among warblers, partridges, and pigeons.




### ATTACK

In other situations, birds adopt openly aggressive behaviors in the presence of intruders or predators. Magpies can even hound and chase away eagles if the latter threaten their brood. Such active defenses are more common among birds of prey.

### PROTECTION

When a parent detects danger, it gets close to its young and covers them so that they are not alone. This behavior is common among tropical birds (tropical seabirds). Several species of curlews and sandpipers place their young between their legs, whereas grebes carry their chicks on their backs when swimming.



**TO DEFEND ITSELF** from the falcon's attack, a flock of starlings squeezes together in a dense formation. If they are near a tree, they do not hesitate to hide in it.





# Diversity and Distribution

**DUCK** (Anatidae family)  
A natural-born fisher, ducks feed on small snails and aquatic insect larvae.

**ONE BIRD, ONE NAME** 64-65

**WHERE THEY LIVE** 66-67

**NO FLYING ALLOWED** 68-69

**MARINE RESIDENTS** 70-71

**FRESHWATER BIRDS** 72-73

**ARMED TO HUNT** 74-75

**TALKATIVE AND COLORFUL** 76-77

**THE PERCHERS CLUB** 78-79



The environment in which an organism usually lives is called its habitat. In their habitats, birds find food, the best places to build nests, and escape

routes in case of danger. An almost universal pattern of distribution shows that more species live in the tropics than elsewhere. With evolution, birds with a common origin have diversified as they

have begun to occupy different environments. This phenomenon is called adaptive radiation. We find ocean birds, which have undergone many changes in order to live near the sea, as

well as birds that live in freshwater environments, in forests, and so on. Each type has acquired special physical traits and behaviors as a result of the adaptive process. ●



# One Bird, One Name


To learn more about different birds, we give each species a name. Ancient peoples grouped birds according to practical traits and mystical beliefs. They used birds as food or considered them to be bad omens or symbols of good luck. The people who developed scientific thought created a classification system that took into consideration the external form as well as the behavior of these vertebrates; hence, the denominations predator, wading bird, and songbird were developed, among others. The most recent system of classification, which is based on genetic and evolutionary criteria, has generated a hierarchical organization of names that is constantly being updated. ●



**HOATZIN**  
*Opisthocomus hoazin*  
is a tropical bird species that inhabits the Amazon. The presence of talons on the chicks' heels links them to their earliest ancestors, including Archaeopteryx.

## Diversity and the Environment

Living birds are distributed among a wide variety of habitats. They can be found in aquatic (freshwater or marine) and aero-terrestrial environments. Marine birds live on cliffs, on islands, or in mangrove swamps. They are excellent fishers, and they use seashores or crannies between rocks as refuges for nesting. In freshwater bodies, such as rivers and streams, ducks feed on plants and surface microorganisms. Muddy shores are rich in insects and mollusks, which are the favorite dishes of ibis. Herons, storks, and egrets spear fish with their sharp bills as they wade in water with their long legs without getting wet. Forests, jungles, mountain ranges, and wide plains form most of the world's aero-terrestrial environments. In jungles and forests, predatory birds hunt their prey, while trogons and parrots gorge on insects and fruit. Rocky peaks are the refuge of condors, which fly for hours in search of the remains of dead animals. Ostriches run over prairies and savannas.



**FALCONIFORMES**  
295 SPECIES  
Condors, Buzzards, Eagles, Vultures, and Falcons

**COLIIFORMES**  
6 SPECIES  
Mousebirds

**PSITTACIFORMES**  
360 SPECIES  
Parrots, Parakeets, Lorries, Cockatoos, and Macaws

**CORACIIFORMES**  
204 SPECIES  
Common Kingfishers and Bee-Eaters

**PICIFORMES**  
382 SPECIES  
Woodpeckers, Toucans, and Puffbirds

**COLUMBIFORMES**  
317 SPECIES  
Pigeons and Doves

**CUCULIFORMES**  
160 SPECIES  
Cuckoos, Turacos, and Hoatzins

**PASSERIFORMES**  
5,400 SPECIES  
Perching Birds and Songbirds

**CHARADRIIFORMES**  
350 SPECIES  
Seagulls, Lapwings, and Plovers

**UPUIFORMES**  
1 SPECIES  
Hoopoe

**STRIGIFORMES**  
174 SPECIES  
Owls

**TROGONIFORMES**  
39 SPECIES  
Trogons and Quetzals

**CICONIIFORMES**  
120 SPECIES  
Herons, Storks, Ibises, and Egrets

**GRUIFORMES**  
190 SPECIES  
Moorhens, Cranes, and Coots

**STRUTHIONIFORMES**  
1 SPECIES  
Ostrich

**PHENICOPTERIFORMES**  
1 SPECIES  
Flamingos

**GALLIFORMES**  
274 SPECIES  
Chickens, Turkeys, Quails, Pheasants, and Partridges

**TINAMIFORMES**  
47 SPECIES  
Tinamous


**CASUARIIFORMES**  
4 SPECIES  
Cassowaries and Emus

**APTERYGIFORMES**  
4 SPECIES  
Kiwis

**RHEIFORMES**  
2 SPECIES  
Rheas

## What Is a Classification?

Since the early Renaissance in the 16th century, scientific classifications assigned compound names to birds, as well as to other living creatures. The first part stands for a genus and the second for a specific name. Thus, the rock (domestic) pigeon is called *Columba livia* in scientific terms. The discovery of new species in different parts of the world widened the array of birds so that compound names were no longer enough. The level of family was established to group genera of birds that share similar traits. In turn, bird families that share common traits were grouped into orders. These were then combined into a category called class. Classes include all current and extinct birds. Birds share a higher category in the hierarchy—the phylum—with fish, amphibians, reptiles, mammals, and vertebrates in general.



**SPHENISCIFORMES**  
18 SPECIES  
Penguins

**GAVIIFORMES**  
5 SPECIES  
Loons (diving birds)


**PROCELLARIIFORMES**  
110 SPECIES  
Albatrosses, Petrels, and Fulmars

**PODICIPEDIFORMES**  
21 SPECIES  
Grebes

**ANSERIFORMES**  
150 SPECIES  
Ducks, Geese, and Swans

**PELECANIFORMES**  
62 SPECIES  
Pelicans, Boobies, and Cormorants

**CAPRIMULGIFORMES**  
109 SPECIES  
Potoos, Frogmouths, and Nightjars





# Where They Live


With their mobility, birds have conquered all areas of the Earth. Despite this characteristic, there are few cosmopolitan species—that is, most birds have specific habitats determined by climate and geographic features. Count de Buffon in the 18th century was the first person to notice that living beings are not distributed homogeneously. By analyzing how animals were dispersed on the planet, he realized that different places had different types of fauna. After the work of naturalist Charles Darwin and ornithologist Philip Sclater, it became clear that organisms are situated in specific biogeographic regions. ●

## Biodiversity in the World

The most diverse regions in terms of bird populations are the tropics because of the favorable conditions of abundant food and warm climate found in them. Temperate regions, however, with their seasons, are destinations for migrating birds from tropical and polar regions. Cold regions, on the other hand, have little diversity but are rich in population density. The rule is that diversity of life-forms happens in places where the environment requires less severe adaptations.


### NUMBER OF SPECIES

- up to 200
- 200 - 400
- 400 - 600
- 600 - 800
- 800 - 1000
- 1000 - 1200
- 1200 - 1400
- 1400 - 1600
- 1600 - 1800



### Nearctic

**732** Species  
62 Families



#### CHARACTERISTICS

- Climatic barrier of cold weather and oceanic isolation
- Most migrating species
- Many insectivorous and aquatic birds
- Affinity with Palearctic

Endemic Avifauna: loons and puffins

Pacific Ocean


NORTH AMERICA



PUFFIN  
*Fratercula artica*

### Oceania

**187** Species  
15 Families



#### CHARACTERISTICS

- Large area and number of climates
- Gliders, divers, and swimmers
- Abundance of fish-eating species
- Many cosmopolitan species


Endemic Avifauna: albatrosses, sheathbills, petrels, penguins, and seagulls

Atlantic Ocean

CENTRAL AMERICA

### Neotropic

**3,370** Species  
86 Families



#### CHARACTERISTICS

- Long-lasting geographic isolation
- Many primitive species
- Great numbers of frugivores

Endemic Avifauna: rheas, tinamous, oilbirds, hoatzins, cotingas, and stripe-backed antbirds

This region undoubtedly has the greatest diversity of birds. The variety in the South American tropics, the most important tropical zone in the world, is one and a half times greater than that of tropical Africa. With more than 1,700 species, Colombia, Brazil, and Peru are the countries with the greatest diversity of avifauna. Even Ecuador, a much smaller country, has more than 1,500 species.

#### HOATZIN

*Opisthocomus hoazin*


SOUTH AMERICA



OSTRICH  
*Struthio camelus*

### Palearctic

**937** Species  
73 Families



#### CHARACTERISTICS

- Climatic barrier of cold weather and oceanic isolation
- Low diversity of species
- Most are migratory species
- Many insectivorous and aquatic birds


Endemic Avifauna: wood grouse, waxwings, flycatchers, cranes

EUROPA

AFRICA

### Afrotropic

**1,950** Species  
73 Families



#### CHARACTERISTICS


- Maritime and desert isolation
- Great number of Passeriformes
- Many flightless birds

Endemic Avifauna: ostriches, turacos, cuckoos

Indian Ocean

### Indomalaya

**1,700** Species  
66 Families



#### CHARACTERISTICS

- Affinities with the Afrotropical zone
- Tropical birds
- Many frugivores


Endemic Avifauna: ioras, pittas, swifts

Pacific Ocean

OCEANIA

### Australasia

**1,590** Species  
64 Families



#### CHARACTERISTICS

- Long isolation
- Many flightless and primitive birds

Endemic Avifauna: emus, kiwis, cockatoos, birds of paradise

RUBY-THROATED HUMMINGBIRD  
*Archilochus colubris*



### COUNTRIES WITH THE MOST SPECIES

More than 1,500

- Colombia
- Brazil
- Peru
- Ecuador
- Indonesia

More than 1,000

- Bolivia
- Venezuela
- China
- India
- Mexico
- Democratic Republic of the Congo
- Tanzania
- Kenya
- Argentina

## Adaptations According to the Environment

Birds are found in all habitats of the world, although most live in tropical regions. Their ability to adapt, however, is remarkable. From jungles to deserts, from mountains to coasts, and even on the sea, birds have succeeded in acclimating themselves. They have undergone a highly varied array of changes in form and behavior. Emperor penguins not only nest in Antarctica but they also incubate their eggs between their feet for 62 to 66 days. The male Lichtenstein's sandgrouse has developed a sponge of feathers to bring water to its chicks, and hummingbirds have special wings that enable them to make all sorts of maneuvers.






# No Flying Allowed

A few birds have lost their ability to fly. Their main characteristic is wing loss or reduction, although for some a remarkable size may be the cause of their inability to fly. Such birds weigh more than 39 pounds (18 kg). This is the case with runners (ostriches, cassowaries, emus, rheas, kiwis), extremely fast birds that live in remote areas of New Zealand, and swimmers, such as penguins, that have developed extraordinary aquatic abilities. ●

## Super Swimmers

Penguins' bodies are covered with three layers of small, overlapping feathers. A penguin has small limbs and a hydrodynamic shape that helps it swim with agility and speed. Dense, waterproof plumage and a layer of fat insulate the bird from the low temperatures of the regions where it lives. Since its bones are rigid and compact, it is able to submerge itself easily. This adaptation distinguishes it from flying birds, whose bones are light and hollow.



**FLIPPERS**  
The short, compact wings look like flippers. They are essential to the penguin's underwater movements.

Rockhopper Penguin  
*Eudyptes crestatus*

## PENGUIN HEADING TO THE WATER



### HUNTING


The wings work like flippers. The foot—with four joined toes pointing backward—and the tail steer the direction of the dive.

### BREATHING

When looking for food, penguins need to leave the water and take a breath between plunges.

### RELAXING

When resting in the water, they move slowly. They float on the surface with their heads up and balance their bodies with their wings and feet.




**Runner's Chest**  
The keel-shaped sternum of flying and swimming birds offers a larger surface for attachment of the pectoral muscles. The flat sternum of running birds has a smaller surface and, consequently, less mobility.


**AFRICAN OSTRICH**  
A single species inhabits eastern and southern Africa. Adults reach a height of 9 feet (2.75 m) and a weight of 330 pounds (150 kg).

## The Ratites

Running birds belong to the group of the ratites (*rata* = raft, an allusion to the flat sternum). The front limbs either are atrophied or have functions unrelated to flying. The hind limbs have very strong muscles as well as sturdy, vigorous bones. Another difference is found in the sternum. It is a flat bone without a keel, which flying and swimming birds possess. Wild ratites can be found only in the Southern Hemisphere. The Tinamidae, native to Central and South America, belong to this group (partridges).




**STRUTHIONIFORMES**  
The ostrich is the only species in this group. It uses its wings for balance when running fast. It has only two toes on each foot. The adult male can weigh up to 330 pounds (150 kg).




**RHEIFORMES**  
Rhea are common in South American countries, such as Argentina. They look like ostriches but are smaller. Their three-toed feet allow them to chase prey. Their long necks and excellent eyesight make them skillful hunters.



**CASUARIIFORMES**  
Agile runners and swimmers. The colors on their necks and heads are distinctive. A bony hoof protects them from vegetation when they run. They have long, sharp talons on their feet.



**APTERYGIIFORMES**  
Kiwis. These birds have four toes on each foot, and their feathers look like fur because they do not have barbules. They normally use their keen sense of smell to find insects at night. They lay only one large egg.



**GREATER DIVERSITY**  
In many cases, running birds can be found in many parts of the world because of human intervention. The area where flightless birds have diversified the most is Oceania, due to continental isolation.

## Running and Kicking

Ostriches usually run to escape from predators or to hunt small lizards and rodents. In both cases, because of their strong legs, they are able to reach a speed of 45 miles per hour (72 km/h) and to maintain it for 20 minutes. When running is not enough to protect the bird, kicking is a valid recourse that discourages the attacker. In courtship displays, forceful stamping is also used to win over females.

18

**VERTEBRAE**  
THE NUMBER AN OSTRICH HAS IN ITS NECK

### ON TWO TOES

With just two toes, the contact surface between the foot and the ground is relatively small. This is an advantage when moving on land.



## Other Walkers

More than 260 species belong to the order Galliformes, which includes chickens, turkeys, and pheasants. The birds in this group have keels, and they perform abrupt and fast flights, but only in extreme situations. Their feet are suitable for walking, running, and scratching the ground. This group includes the birds that human beings use the most. In general, males are in charge of incubating and raising the young.

### FLYING WITH LITTLE GRACE







# Marine Residents

Of the more than 10,000 bird species inhabiting the Earth, only about 300 have managed to adapt to marine life. To survive at sea, they have undergone multiple adaptations. For instance, marine birds have a more efficient excretory system than that of other species, including a specific gland that helps them eliminate excess salt. Most marine birds live on the coasts and have mixed behaviors; others are more aquatic than aerial. A few—such as albatrosses, petrels, and shearwaters—can fly for months at a time, landing only to raise their chicks. They are called pelagic birds.

## Adaptations

Marine birds are well prepared to live on water, especially those that fish out at sea. The tips of their bills are hook-shaped, and their feet have webbed membranes between the toes. They also have an admirable ability to float. The saline water is


not a problem; these birds can even drink it. In some pelagic birds, sense of smell plays an important role in enabling them to detect the oil of the fish in the water to find schools of fish. They also use their sense of smell to find their nests in colonies.



**TOTIPALMATE FEET**  
A characteristic of many marine birds. The posterior toe (hallux) is joined to the other toes by a membrane. It creates more surface area, and therefore more push, as the bird swims. Birds with this kind of foot walk clumsily.

## VARIOUS MARINE BIRDS

**IMPERIAL SHAG**  
*Phalacrocorax atriceps*  
This great coastal diver has solid bones and strong swimming feet. It does not oil its plumage so that it can better submerge.



**BROWN PELICAN**  
*Pelecanus occidentalis*  
It stays on the shore. It uses its crop as a fishing net while it swims.



**HERRING GULL**  
*Larus argentatus*  
A voracious fisher and great glider. There are many species, some truly cosmopolitan.



**CAPE GANNET**  
*Morus capensis*  
Skillful spearfishers. They live in colonies in Africa. To help cool themselves, they have a strip of naked skin on their throats.



148 feet (45 m)

IS THE MAXIMUM DEPTH THEY REACH. COMMON LOONS—BIRDS INDIGENOUS TO NORTH AMERICA THAT SPEND THE WINTER AT SEA—HAVE BEEN RECORDED TO REACH THIS DEPTH. ALMOST UNABLE TO WALK, COMMON LOONS ARE EXCELLENT SWIMMERS AND DIVERS. THEY NEST AT INLAND LAKES DURING THE SUMMER.



**COMMON LOON**  
*Gavia immer*


**HOOK-SHAPED TIP**  
Prevents the fish from sliding and falling away, once caught

**TUBULAR NOSTRILS**  
Albatrosses have one on each side of their bills. On petrels and shearwaters, the tubular nostrils have merged on top of the bill, forming a single nasal tube.

**BILL**  
It is composed of several hard plates.

## Salt Gland

Living in the ocean requires a few adaptations. The most notable one is the salt gland, which eliminates excess salt from the bloodstream. This way, marine birds can even drink salt water without suffering dehydration, as would be the case with humans. This gland is very efficient: it has been observed that 20 to 30 minutes after drinking a saline solution with concentrations similar to that of the ocean (4 percent), birds eliminate another solution (through the nostrils) with 5 percent salt, in the shape of water drops.



## Fishing Methods

Many marine birds fish by diving into the sea. This way, they can access fish that swim below the surface. In order to reach deeper levels in the water, they fly up several feet, spy a school, fold their wings, and plunge with their necks stretched forward. Thanks to the buoyancy of their feathers, they are back on the surface moments later.



## DIVE FISHING

1. The bird dives to gain speed.

2. It folds its wings and stretches out its neck to immerse itself in the water and reach the school of fish.

3. The bird immerses its body as much as possible to catch the fish; its feathers cause it to float back up.




# Freshwater Birds

This group includes birds that vary greatly—from common kingfishers to ducks to storks—and covers a wide spectrum. Freshwater birds live in rivers, lakes, and ponds for at least part of the year and are perfectly adapted to aquatic life. Some are excellent swimmers, whereas others are great divers. An important group wades in watercourses with long legs as they fish. Freshwater birds have a varied diet and are mostly omnivorous. ●

## Ducks and Distant Cousins


The order Anseriformes includes birds that are very familiar to humans: ducks, geese, and swans, for example. They have short, webbed feet and wide, flat bills lined with lamellae (false teeth) that enable them to filter their food, catch fish, and scrape the beds of rivers and ponds. Most are omnivorous and aquatic (either staying on the surface or diving), although some species spend more time on land. They are widely distributed, and the plumage of males becomes very colorful during the courtship season.




Fulvous Whistling Duck  
*Dendrocygna bicolor*

### HOW THEY USE THEIR FEET TO SWIM

A duck moves its feet in two ways. To advance, it spreads out its toes and uses its webbed feet to row. It closes the toes to bring the foot forward again. If the bird wants to turn, only one foot pushes to the side.



### A DUCK'S DIET




**ORIFICES**  
Open and oval

**LAMELLAE**  
Around the inside edges of the bill

### DUCK BILLS

are flat, wide, and slightly depressed toward the middle. In general, their shape does not vary, but there are species with tiny bills (the mandarin duck, for example).




**Shovel-Shaped Bill:** Typical of many ducks. The size varies.

**Mandarin Duck Bill:** One of the smallest-billed species.

## Divers and Other Fishers


Diving birds belong to the family Podicipediformes. They feed on small fish and aquatic insects. They are very clumsy on the ground. In the Coraciiformes order, common kingfishers and other similar birds find their prey by closely watching the water. When one of these birds notices a small fish, it spears it, catching it with its bill. In the order Charadriiformes, curlews wander around the edges of ponds in search of food. Their long legs keep their bodies out of the water. They are not swimmers.



**THE BILL OF AN IBIS** is long and thin, ideal to stick in the mud to look for food.

## Wading Birds

These birds belong to an artificial order since, from a genetic perspective, the species are not related. They are grouped together because adaptation to the same habitat has caused them to develop similar shapes: long bills and necks to perform skillful movements and thin legs designed to wade across the water as they fish. Herons form a special group because they are cosmopolitan and because they have powder down feathers. Ibis and storks also have a wide distribution (area in which they occur). Birds that have spoon- and hammer-shaped bills are found primarily in Africa.



White Ibis  
*Edocimus albus*

**THE LEGS OF AN IBIS** keep the bird above the water but close enough to fish. Ibis also stir up the beds of lakes and ponds.



# Armed to Hunt

**B**irds of prey are hunters and are carnivorous by nature. They are perfectly equipped to eat living animals. Their eyesight is three times sharper than that of human beings; their ears are designed to determine the precise status of their prey; they have strong, sharp talons; and they can kill a small mammal with the pressure of their talons alone. Their hook-shaped bills can kill prey by tearing its neck with a single peck. Eagles, falcons, vultures, and owls are examples of birds of prey. Birds of prey can be diurnal or nocturnal, and they are always on the lookout. ●

## Diurnal and Nocturnal

Eagles, falcons, and vultures are diurnal birds of prey, whereas owls are nocturnal—that is, they are active during the night. These two groups are not closely related. These birds' main prey includes small mammals, reptiles, and insects. Once they locate the victim, they glide toward it. Nocturnal birds of prey are specially adapted: their eyesight is highly developed, their eyes are oriented forward, and their hearing is sharp. The feathers on their wings are arranged in such a way that they make no noise when the bird is flying. In order to protect themselves while sleeping during the day, they have dull plumage, which helps them blend in with their surroundings.



**EURASIAN EAGLE OWL**  
*Bubo bubo*  
Its ears are asymmetrical and can determine the location of prey with great precision.



**BALD EAGLE**  
*Haliaeetus leucocephalus*  
It has a visual field of 220 degrees and a bifocal vision of 50 degrees.

**CERE**  
Fleshy formation, somewhat thick and soft

## Bills

The bills of birds of prey are hook-shaped. Some birds of prey have a tooth that works like a knife, allowing them to kill their prey, tear its skin and muscle tissues, and get to the food easily. The structure and shape of the bills of birds of prey changes depending on the species. Scavengers (for example, vultures and condors) have weaker bills because the tissues of animals in decomposition are softer. Other species, such as falcons, catch prey with their talons and use their bills to finish it off with a violent stab to the neck, breaking its spine.

**TIP**  
Where the tooth is located

**NOSTRIL**  
Olfactory canals

**Zone-Tailed Hawk**  
*Buteo albonotatus*



**BALD EAGLE**  
Its hook-shaped bill is common to many birds of prey.



**SPARROW HAWK**  
Its thin bill enables it to take snails out of their shells.



**FALCON**  
It can break the spine of its prey with its upper bill.



**GOSHAWK**  
Its strong bill can catch prey as large as hares.

## OWL PELLETS

Owls produce pellets. They swallow their prey whole and regurgitate the indigestible substances. The study of pellets makes it possible to determine the fauna of small areas with great precision.



**5 miles**  
(8 km)

IS THE DISTANCE FROM WHICH A FALCON CAN PERCEIVE A PIGEON.



**GRIFFON VULTURE**  
Its long toes do not have a good grasp.



**FISHING EAGLE**  
Its toes have rough scales that look like thorns, which help it to catch fish.



**GOSHAWK**  
It has calluses at the tips of its toes.



**SPARROW HAWK**  
Its feet have tarsi and short, strong toes.

## HOW THE VULTURE HUNTS



1

Vultures feed mainly on carrion, although they are able to attack a living animal if it is vulnerable and the situation presents itself.

2

Thanks to their ability to glide on thermals, vultures can find carcasses on which to feed without wasting energy.

3

Once they find food, they must analyze the territory to know if they will be able to take flight again soon.



## DIMENSIONS

The wings of birds of prey are adapted to suit their flying requirements. They can measure up to 10 feet (3 m).



Condors 3 to 9.5 ft (0.95–2.9 m)



Eagles 4.5 to 8 ft (1.35–2.45 m)



Buzzards 4 to 5 ft (1.2–1.5 m)



Kites 2.6 to 6.4 ft (0.8–1.95 m)



Red-Backed Hawk 3.4 to 4.4 ft (1.05–1.35 m)



Falcons 2.2 to 4.1 ft (0.67–1.25 m)

## Feet

Most birds of prey catch and kill their prey with their talons and tear away the meat with their bills. For this reason, birds' feet constitute one of the morphological characteristics of a species. The toes end in strong, sharp nails

that the bird uses as pincers to catch its prey in flight. The osprey also has thorns on the soles of its feet, which help it to catch fish.




# Talkative and Colorful

Parrots form a very attractive bird group with a great capacity for learning. This group comprises cockatoos, macaws, and parakeets. They share physical characteristics, such as a big head, a short neck, a strong hook-shaped bill, and climbing feet. They have plumage in many colors. Toucans and woodpeckers share with parrots the colors of their feathers and their type of feet. Toucans have a wider, thicker bill, but it is light. Woodpeckers are climbing birds with a strong, straight bill, a tail of stiff feathers, and a distinctive crest. They form numerous groups, and most nest in trees. ●

## Eating, Climbing, and Chattering


Parrots use their bills to feed and to move about tree branches; they use their bills as an extension of their feet to give them support when they climb. Parrots also have a curved profile, a pointed tip on their upper mandible, and sharp edges on their lower one. These adaptations are practical when it comes to cutting and breaking fruits and seeds. The largest species like fruits with shells, such as walnuts, hazelnuts, and peanuts. The smaller ones prefer nectar and pollen, which they obtain with the help of brush-shaped bristles on their tongues. Their ability to imitate the human voice has made them very popular. However, they are far from being able to produce language. In reality, they are merely good imitators: they use their excellent memory to imitate sounds. They do this when they are hungry or when they detect the presence of unknown people.



**WINGS**  
Usually they are short and rounded, suitable for flying among branches.

**NOSTRILS**  
They are located at the base of the bill's upper portion.

**HOOKEED BILL**



**The Feet**  
are referred to as zygodactyl. This means that two toes project forward and two project backward. Parrots appear to strut because their feet have a tibiotarsus that is shorter than that of other birds.



**FEET LIKE HANDS**  
In some species, the left foot is longer. They use it to grab fruits and tear them with their bills.

**FEATHERS AND COLORS**  
They have tough and lustrous plumage. An abundance of green feathers helps them to hide among the leaves. In South America, the array of colors includes hues of blue, yellow, and red.



**WOODPECKERS**  
hollow out tree trunks with pecks in order to build a nest and to feed on insects that eat wood.

### THE HABITAT OF WOODPECKERS

They live in the woods and can often be heard there. Their adaptations to arboreal life are demonstrated by their strong, thick bills and their stiff tails, which they use for support, together with their feet. They use their hearing to locate tree-boring insects; they then peck the wood incessantly until they find them.



**TOUCANS**  
Their big bills have serrated edges that suit their diet of fruit. They live in the South American jungles.

**QUETZALS**  
They belong to the family Trogonidae. They have feet adapted to arboreal life. Males have brilliant plumage and long, attractive tails.






# The Perchers Club

Passerines—or Passeriformes, the scientific name—form the widest and most diverse order of birds. What distinguishes them? Their feet are suited for perching and, therefore, for living among trees, although they can also stroll on the ground and through the brush. They inhabit terrestrial environments all over the world, from deserts to groves. Their complex sounds and songs originate from a very well-developed syrinx. Their chicks are nidicolous—that is, naked and blind at birth. In their youth, they are agile and vivacious, with very attractive, abundant, and colorful plumage. ●

## THE SMALLEST

Passerines are small in comparison with other birds. Their size varies from 2 inches (5 cm) (bee hummingbirds, *Mellisuga helenae*) to 7 inches (19 cm) (*Chilean swallow, Tachycineta leucopyga*) to 26 inches (65 cm) (common raven, *Corvus corax*).



**HUMMINGBIRDS** 2 in (5 cm)  
They get so much energy from nectar that they can double their body weight by eating. However, they use this energy up during their frantic flights.

**SWALLOWS** 7 in (19 cm)  
Swallows have great agility and skill. These popular migratory birds have bodies suited for long trips.

**RAVENS** 26 in (65 cm)  
They eat everything: fruits, insects, reptiles, small mammals, and birds. They are skillful robbers of all kinds of food.

## PASSERIFORMES BIRDS

Passerines have been classified into 79 families, with more than 5,400 different species.



## SINGER


This blue-and-white swallow (*Notiochelidon cyanoleuca*) intones its pleasant and trilling chant while it flies or when it alights. Larks, goldfinches, canaries, and other passerines delight us with their trills and sounds.

## HARD, SHORT BILL

The bill of a swallow is very short and tough. The swallow can use it to catch insects in flight.


## SYRINX

This sound-producing organ is located at the end of the trachea. The muscles in the syrinx move the bronchial walls, which, as the air passes through, produce the melodic sounds that characterize songbirds.



## LIVING AT THE EXTREMES

They range from one hemisphere to the other. They raise their chicks in the north and fly to the south to spend winter there. They fly all the way to Tierra del Fuego. Their sense of direction is remarkable. They can find and reuse their nests after returning from a migration.



**A** In the summer, during the reproductive season, they live in the Northern Hemisphere on the American continent. In general, neotropical migratory birds are those that reproduce above the Tropic of Cancer.

**B** When winter arrives in the Northern Hemisphere, they perform a mass migration to the south, occupying the Caribbean and South America. The barn swallow travels 14,000 miles (22,000 km) during its migratory trip from the United States to southern Argentina.

## PERCHING FOOT

Three toes project forward, and the well-developed hallux projects backward. This type of foot allows the bird to hold on tightly to branches.

## BARN SWALLOW

(*Hirundo rustica*)  
Barn swallows spend most of their time traveling to temperate zones.

## Family Album

Four basic groups have been established to facilitate the study of families: passerines with wide bills; ovenbirds, whose plumage is dull and brown (ovenbirds are noted for the great care they take in building nests); lyrebirds, whose tails have two external feathers that are longer than the others; and songbirds, with their elaborate and pleasant singing. Songbirds form the most numerous and varied group; it includes swallows, goldfinches, canaries, vireos, and ravens.



## LYREBIRDS

There are only two species of these Passeriformes, and they are found only in Australia. They are very melodic and are excellent imitators of other birds. They can even imitate the sound of inanimate objects, such as horses' hooves.



## WIDE BILLS

They are native to Africa and Asia and inhabit tropical zones with dense vegetation. They eat insects and fruits. They produce nonvocal sounds with the flapping of their wings. They do this during courtship, and the sound can be heard 200 feet (60 m) away.



## OVENBIRDS AND THEIR RELATIVES

Their nests are completely covered structures, similar to ovens. Other members of this family build nests with leaves and straw, weaving interesting baskets. Still others dig tunnels in the ground.



# Humans and Birds

**ROBIN CHICKS** (*Erithacus rubecula*)  
Although their natural habitats are humid groves, they usually seek shelter in cities, always close to water.

**BIRDS AND HUMAN CULTURE** 82-83   **BIRD DOMESTICATORS** 88-89  
**HOW TO GET TO KNOW BIRDS** 84-85   **ENDANGERED SPECIES** 90-91  
**AMONG US** 86-87



**H**uman beings have long taken an interest in studying these high-flying creatures. They have served as a source of food and have sometimes been

used as indicators of the arrival of rain, storms, or the presence of common enemies, such as dangerous reptiles. Evidence of ancient peoples' veneration of birds can be found in documents,

paintings, and reliefs. The Egyptians were the first to domesticate pigeons. Today several species brighten up homes with their colors and cooing. Other types of birds, such as sparrows

and swallows, live with us in cities. The destruction of bird habitats, through the excessive exploitation of natural resources, is one of the main causes of bird extinctions. ●



# Birds and Human Culture

Birds fly, sing, dance, and have showy plumage. Because of these qualities, they have fascinated human beings throughout history. Some species, such as eagles, have played an important role in world literature because of their aggressiveness and beauty. Some birds have also been assigned symbolic meanings: doves, which currently represent peace, are one example. Human beings have also been able to make use of birds. In the past, they were often used by sailors to find land, and in other cases they were trained to hunt. ●

## Rites and Beliefs

Birds have long enjoyed a prominent place in religion—first as totems and then as iconic representations of gods. Many religions have featured deities with the wings or heads of birds. Birds were also celestial messengers, and the future was interpreted through their flights. The crow was Apollo's messenger in ancient Greece; the Maya and Aztecs had Quetzalcóatl, their supreme god, who was named after the quetzal (a brightly colored Central American bird); and the Egyptians represented their fundamental god Horus with a falcon.

**HORUS, THE FALCON**, is an important god in Egyptian mythology. His eyes represent the sun and the moon, and together with Seth he watches over the boat of Ra, which carries the dead away on the Nile.

## ELEMENTS

Birds and masters wear specific clothes. In addition to gloves, hoods, and straps, radio transmitters are now also used to locate the birds as they fly.



### FENGHUANG

was the messenger bird of Xi Wangmu, goddess of fertility and eternity in ancient China. A detail of a painting in the caves of Mogao, Dunhuang, China, is pictured above.



### QUETZAL

Pictured above is a detail of a Mayan ceramic piece featuring quetzals, Mesoamerican birds with long green tails, from which, according to myth, the god Quetzalcóatl took his clothing and name: "feathered serpent."



### GARUDA FRESCO

Pictured above is a winged deity, featured on a fragment of a mural from the Hindu temple of Garuda, in Ananta Samarkhom, Bangkok, Thailand.

## Bird Symbolism

Throughout history and across cultures, human beings have used birds to symbolize several concepts. The fascination that their flight creates was a source of inspiration for such interpretations. Today the strongest and most widespread association is that of the flight of birds with freedom. In distant times (and in not so distant times), however, birds have represented many other things, from fertility and happiness, with their spring songs, to deep mourning, in the case of crows and vultures. Wisdom has been associated with owls, and shrewdness with crows. According to a certain modern tale, storks are responsible for bringing babies, and eggs are the universal symbol of gestation.

### EAGLE

In Greek mythology, it was the symbol of Zeus. The Romans used it on their legions' banners. For many native North American cultures, it represented war, and it was the emblem of feudal lords and emperors. Today it is the national symbol of Mexico and the United States.

### DOVE


Doves currently stand for peace, but in ancient Greece, Syria, and Phoenicia, they were used as oracles. In Mesopotamia and Babylon, they embodied fertility. For Christians, they symbolize the Holy Spirit and the Virgin Mary.

## Dressing Up in Feathers

Almost all cultures have used bird feathers for decorative and ritualistic purposes. Their use as ornaments extended to North and South America, Africa, and the Western Pacific. North American indigenous peoples featured them on their war outfits, Hawaiian kings wore them on their royal costumes, and the Mayans and Aztecs used them in works of art.

## Falconry

This practice originated in Asia, in the homeland of the nomadic Mongols—descendants of Genghis Khan—where, to this day, it is commonly used as a form of subsistence by part of the population. It consists of using birds of prey (mainly falcons) to hunt. Trained birds are typically carried, hooded, and perched on their masters' arms. When released, they fly at high altitudes looking for prey, and then dive toward the ground to hunt it. They carry the prey back to their masters, who reward them with food. The basic training process takes a little over a month and a half.



HOOD

FALCON

TRANSMITTER

GLOVE

LASH

STRAPS


**PIGEON** Sometimes their numbers in urban areas become excessive.

### COMPETITOR BIRDS

When birds share a habitat with humans, they often compete for resources (light, water, space, and nutrients). This is the case with birds that feed on cultivated crops. Urban areas, which have buildings that offer good nesting sites, attract many birds. This fact can be commonly observed in squares and open spaces, where pigeons and sparrows form veritable flocks.

**SPARROW** The sparrow is one of the birds best adapted to the urban environment.

**NORTH AMERICAN INDIGENOUS MAN** wearing a war costume covered with feathers





# How to Get to Know Birds

**O**rnithology (from the Greek *ornitho*, "bird," and *logos*, "science") is a branch of zoology that studies birds. Ornithologists and a great number of bird lovers, who enthusiastically want to know more about these creatures, carry out the task methodically and patiently. They observe, analyze, and register birds' sounds, colors, movements, and behavior in their natural environments. To undertake this fieldwork, they develop methods and techniques and use technological resources to track specimens and learn about what happens to them during given times of the year. ●

## WORK CLOTHES

Although seemingly a small detail, clothes can be a hindrance. They should be comfortable and soft, appropriate for the weather, and of colors that blend into the environment.

## BINOCULARS

They make it possible to see details in color and shape without disturbing the birds. Their usefulness depends on the power of their lenses.

## Direct Observation

Observing in a bird's natural habitat provides much information. For the greatest success, bird-watchers typically place themselves in front of rocks or trees in order not to form a silhouette. An alternative technique is to create a hiding place, such as a hollow cardboard rock. In both cases, the watcher needs to be facing away from the sun and must be prepared to stay for a long time.

## STUDIES

Many studies on anatomy, physiology, and genetics of different bird species are carried out in laboratories.

## Catching Birds


### MIST NET

These fine nets are usually placed over swamps or marshes, and they can catch small birds. Once the birds are identified with a ring or some other tag, they are set free.




### CANNON NETS

These nets are "shot" over birds using cannons or rockets. They unroll and catch birds as they are eating or resting. They are used to catch large birds.



### HELIGOLAND TRAP

It consists of a large barbed-wire funnel or corridor that ends in a box. Birds caught like this are tagged so that they can be monitored and studied later on.




### CAMERA WITH ZOOM LENS

A camera with a powerful lens provides the opportunity to record details that simple cameras cannot.

### TAPE RECORDER

Recording birds' noises and songs makes it possible to distinguish among birds. Experts can identify species by listening to their recordings.



## The Marking of Captured Birds

This technique provides data on migration, survival, and reproduction rates, among other data. The bird should not suffer adverse effects in its behavior, longevity, or social interactions. Under no circumstances should this procedure hurt a bird. To avoid hurting birds, rings are designed to be placed on them rapidly and easily, yet to stay in place until the research is completed.

### RINGED


Numbered aluminum rings are used. When placing one on a bird, one should make sure that it slides and turns around the tibiotarsus to avoid hurting the bird or causing it to change its activity.

### WING MARKERS

They are very visible and can be codified for individual identification. They stay on the bird for long periods of time and are normally used on birds of prey.


### NECK MARKERS

When placed appropriately, necklaces are effective markers with few adverse effects on geese, swans, or other aquatic birds with long necks.



### NASAL MARKERS

These are colored, numbered plastic disks placed on the bill. They are fastened to the nasal orifices of aquatic birds.





### PAINTS AND DYES

Birds that visit environments with dense vegetation are normally marked with nontoxic colorings on the feathers of their highest and most visible body parts.



Scientists take advantage of molting to study penguins. When they are changing plumage, these birds stand on their legs, which makes it difficult to place or observe any rings on their legs. Instead, strips are placed around their wings, or electronic chips are implanted in their skin. The latter technique is less harmful because it does not potentially hinder the animal.





# Among Us

The urban environment presents opportunities for birds. It offers advantages in finding food and shelter. People, young and old, give bread crumbs to these interested visitors. When birds come to cities, houses and parks offer them protection. They can find more options when it comes to building a nest. Seagulls and owls have changed their behavior by adapting to the city, and other species, such as some sparrows, are no longer able to survive without a human presence. However, not everything is an advantage. In the city, birds have to face dangers and obstacles that do not exist in their natural habitat: a utility wire or a car can be fatal. ●

## Where to Find Them

In big cities, groups of birds can be found in different areas. Busy and noisy areas, such as squares, parks, and gardens, attract many species. Calm, deserted spaces, such as cemeteries or deserted buildings, are chosen by birds in search of peace. Other places where birds choose to eat and sleep include plots and landfills with lots of food, as well as nooks in high places, such as balconies, belfries, and roosts.

### PLOTS AND CEMETERIES

Plants with seeds usually grow in these places, which are located far from urban centers. For this reason, magpies and nightingales favor them.

### DAY/NIGHT

Artificial night lighting intensifies birds' activities. They organize their tasks according to street lighting.

### PIGEONS

spend a lot of time together. They are granivores but eat all kinds of food.

**SWIFTS** are insectivores. They catch their prey in flight.

### SEAGULLS

usually fight among themselves for the food they get from landfills.

### URBAN CENTERS

With the arrival of cold weather, different bird species from the countryside and the mountains come to the city. In general, they stay until the beginning of spring. In the winter, more birds, such as chiffchaffs, great tits, and robins, can be observed in cities.

### FALCONS

Diurnal birds of prey. They nest on the roofs of high buildings.

### SPARROWS

Small birds with a highly varied diet

**BLACKBIRDS** were originally migratory, but as they adjusted to cities, they became nonmigratory.



## The Urban Environment

This setting is characterized by environmental and climatic factors that are different from the natural ones. It has more varieties of plants, higher average temperatures, less wind, more rain, cloudier skies, and less solar radiation. Polluted air and soil are harmful factors for both humans and birds.

**10%** RAIN INCREASE  
**15%** WIND INTENSITY DECREASE  
**1.5°** TEMPERATURE INCREASE DOWNTOWN

### GROVE

**WAGTAILS** nest in holes or crannies, always near water.

**ROBINS** They get close only to find food. Their trills and plumage are attractive.

**ABANDONED BUILDING**

**FEEDER WITH FRUITS**

### DIET



Some species, which are veritable strategists, benefit from human activities. For example, seagulls can feed on garbage; great tits drink milk from containers by adroitly uncapping them; and some magpies have learned to tear cardboard containers to eat the eggs inside.

## Population Control

Without enough predators and with an abundance of food, urban bird populations grow exponentially.

**800,000** RAVENS, CROWS, AND RELATED BIRDS LIVE IN MOSCOW.

**160,000** PIGEONS DESCEND ON THE SQUARES AND STREETS OF BARCELONA.



**STORKS** build their nests on rooftops.

**LANDFILL**

### OWLS

Belfries or abandoned buildings are the homes or dormitories of these birds.





# Bird Domesticators

The breeding of birds in captivity has great social and economic value. This activity is carried out all over the world on industrial poultry farms and family farms where birds are raised for consumption and sale. A great variety of domestic birds have been developed from species inhabiting natural environments. We use their flesh and eggs as food and their feathers in coats to protect us against the cold. We also use birds for communication and as colorful and melodic pets. They are so dependent on people that in some cases they cannot survive when they are freed. ●

## At Your Service

Domestic birds have been bred from the following orders: Galliformes (hens, quails, turkeys, and pheasants), Anseriformes (ducks and geese), Columbiformes (pigeons), Passeriformes (canaries), and Psittaciformes (parakeets and parrots). In poultry farming, they are divided according to their use: barnyard birds (Galliformes, Anseriformes, and Columbiformes) and companion birds or pets (Passeriformes and Psittaciformes). Commercial poultry farming of barnyard birds generates high revenues worldwide. Farmers can take advantage of the fact that birds are very active during the day, that they readily live in groups, and that they have a high reproductive rate due to polygamous behavior. Pets have commercial appeal, with their colorful plumage, ability to express themselves, and friendliness toward humans. These characteristics make them much-appreciated pets.



**TURKEY**  
On the American continent, these birds were domesticated by the indigenous pre-Columbians from a wild Mexican species called *Meleagris gallopavo gallopavo*.



**GOOSE**  
Contemporary domestic breeds descend from wild Asian and eastern European species. They are voracious, which makes fattening them easy.



**DUCK**  
These birds are an important source of food in Southeast Asia. In Central and South America, the consumption of ducks is less widespread.



**CAGED**  
Canaries, native to the Canary Islands, have been selectively bred by humans for nearly four centuries.

## Farm Model

When compared to other farm animals, birds grow and reproduce easily. They need to have a place with appropriate temperature, humidity, and ventilation in order to yield the desired amount of meat or eggs. For this reason, it is necessary to maintain continuous environmental and sanitary control of the area in which they are bred. Ideally they should be able to walk, run, scratch the earth in search of food, and take sunbaths. Additionally, to protect them from predators and from inclement weather, it is important to shelter them in coops. This allows them to rest peacefully and sleep at night.

**QUENCHING THIRST**  
Ten chickens drink from 0.5 to 0.8 gallon (2-3 l) of water a day. The farmer provides them with water in troughs, which are placed all over the henhouse.

**MIXED DIET**  
Birds look for insects and plant shoots as they peck the soil. The breeder complements this diet with nutritionally balanced foods.



**AIRMAIL**  
For more than 1,700 years, human beings have used pigeons to send messages. Armies have used them as communication aids during wars. Pigeon keeping is the practice of breeding and preparing pigeons to become messengers, a task that makes the most of their agility and intelligence.

## BIRD FLU

This disease, also called avian influenza A, is caused by a virus whose strains have various levels of virulence. It disseminated from Asian markets, where the overcrowding of domestic birds is common. This promoted the spread of the disease to wild birds. As of 2006, more than 30 million birds had succumbed to this disease. Cats, pigs, and human beings have also been infected.

**50%**


**THE COMMOTION AND FEAR OVER DISEASE HAVE REDUCED THE DEMAND FOR THE CONSUMPTION OF DOMESTIC BIRDS IN MAJOR EUROPEAN CITIES BY HALF.**

**PATHS INTO THE BODY**  
Digestive Tract  
Urogenital Tract

Conjunctiva  
Respiratory Tract  
Needles  
Skin  
Wounds

**SIZE COMPARISON**

Bacteria  
Virus



## DOMESTICATION IN HISTORY

The domestication of birds is a very old activity, as shown by records from different cultures in different parts of the world. It was related to the adoption of a sedentary way of life by human populations.

5000 BC	2000 BC	1492
<b>India</b> There are records of domestic geese in India, beginning in 5000 BC.	<b>Far East</b> Descendants of the royal duck ( <i>Anas platyrhynchos</i> ) were domesticated in this area of the Asian continent (what is now China).	<b>Mexico</b> The Spanish colonizers encountered turkeys domesticated by inhabitants of the New World.



# Endangered Species


Since early civilization, people have affected the Earth's environment. The cutting of trees in rainforests and woodlands has destroyed many bird habitats, the loss of which is the leading cause of bird extinctions today. Also, the introduction of animals such as cats, dogs, and rats to new areas has created a threat for many bird species. Indirect poisoning with pesticides, the trafficking of exotic birds as pets, and the sale of feathers have done further damage to many species. Fortunately, all is not lost. The first step to conserving the world's avifauna is to learn about the extinction of birds and its magnitude.

## The Most Important Causes

Birds are very sensitive to changes in their habitats, and this is the main cause of extinction (87 percent of species are affected by it). Excessive hunting is another of the greatest dangers, affecting 29 percent of the endangered species in the world. The introduction of foreign species is yet another major danger, jeopardizing 28 percent of species. In addition, the intervention of human beings through the destruction of habitats and the introduction of pollution combined with the occurrence of natural disasters harms more than 10 percent of species.

### POISONING

Most birds of prey are endangered by the excessive use of nonbiodegradable pesticides.




### SAVING THE PEREGRINE FALCON FROM EXTINCTION

- 1942** There were 350 couples of peregrine falcons in the United States.
- 1960** Peregrine falcons disappeared in the wild because of excessive use of pesticides (DDT and dieldrin).
- 1970** Falcons were bred in captivity at Cornell University, to be set free later.
- 1986** 850 birds were set free in the southern United States.



**CALIFORNIA CONDOR**  
*Gymnogyps californianus*  
Until 1978, there were 30 specimens in the wild. Bred in captivity, new specimens have been set free since 1993. Their adaptation is being studied.



**HYACINTH MACAW**  
*Anodorhynchus hyacinthinus*  
It is estimated that 1,000 to 9,000 specimens live in the Amazon.

**MANGROVE FINCH**  
*Camarhynchus heliobates*  
There are about 100 remaining species on the Galapagos Islands.


**ROYAL CINCLODES**  
*Cinclodes aricomae*  
live on humid mountain ranges, at altitudes between 11,500 and 15,000 feet (3,500-4,500 m). Their number is unknown.



### CLASSIFICATION OF RISK

- Extinct in the Wild** Species surviving only in captivity
- Critical Risk** Extinction is imminent
- Endangered** Fast-decreasing population
- Vulnerable** High risk of extinction in the wild


### BIRDS OF THE WORLD




### EXTINCT BIRDS

Although the responsibility of human beings is undeniable, many species became extinct because of natural phenomena. However, all the extinctions from the 18th century to date are related to human activities.

**129 SPECIES HAVE GONE EXTINCT SINCE THE 18TH CENTURY.**



**Dodo from Mauritius**  
Quickly exterminated by colonizers and seafarers in the 17th century





# Glossary

## Adaptation

Change in the body of a bird or another animal that allows it to reproduce better in a given environment

## Adaptive Radiation

Evolution of an initial species, adapted to a given way of life, into other species, each adapted to its own way of life

## Aerodynamic

Having an appropriate shape to decrease resistance to the air. Birds' bodies are aerodynamic.

## Alulae

Rigid feathers whose function is to decrease air turbulence during flight

## Amino Acid

Molecule from which proteins are produced

## Ancestor

Progenitor, more or less remote, that passes down a set of characteristics to its descendants

## Angle of Attack

The change in position of a bird's wing to increase or decrease speed and altitude while hunting prey by air

## Apterylae

Naked areas of the skin where feathers do not grow

## Atrophy

Significant decrease in the size of an organ. The wings of nonflying birds have undergone atrophy during evolution.

## Barbs

Thin, straight, parallel blades, perpendicular to the shaft. They resemble the leaves of a palm tree.

## Bill

Hard cover of a bird's mandibles; also called the bill.

## Biodiversity

Variety of species that live in a given natural or artificial environment

## Biogeographic Regions

Geographic regions that biologists analyze to determine the distribution of animals and other living organisms, according to the geographic conditions of a place. Migratory birds usually travel through different biogeographic regions between winter and summer.

## Biped

Aero-terrestrial animal that walks on its posterior limbs. Birds are bipeds.

## Briny

Water sample or body of water containing between 0.08 and 4.25 ounces (0.5-30 g) of salt per gallon of aqueous solution

## Bronchus

Each of the branches into which the trachea divides. The syrinx originates in the bronchi.

## Calamus

The lower part of the vane that is wider, hollow, and, in general, naked. The feather is attached to the skin through the calamus.

## Camouflage

A characteristic that enables the animal to blend into its environment. It allows the animal to go unnoticed in the presence of predators.

## Carnivore

An animal that feeds on meat

## Carrion

The remains of dead animals used as food by some birds or other animals. Vultures are scavengers (i.e., animals that feed on carrion).

## Center of Gravity

Point at which the sum of all the gravitational forces that act on a body converge

## Cere

Thin skin layer that covers the base of the bill

## Chick

A baby bird that has just come out of the eggshell and that has not yet left the nest. Its diet and safety depend on its parents.

## Chorion

One of the coverings that wraps the embryo of reptiles, birds, and mammals

## Class

One of the many divisions scientists use to classify animals. Birds form a class of their own.

## Climate

Average temperature, humidity, and pressure that determine the atmospheric conditions of a region and that are related to other geographic characteristics of that region

## Cloaca

The widened and dilatable final portion of the intestine of a bird or other animal in which the urinogenital tubes converge

## Courtship

Behavior patterns that males and females follow to try to attract partners

## Coverts

Layers of contour feathers that provide a bird's body and wings with support and an aerodynamic surface

## Crepuscular

Active at dawn or twilight, when there is little light

## Crest

Extended or raised feathers located on the upper part of a bird's head

## Crop

Membranous sac that communicates with a bird's esophagus, where food is softened

## Dermal Papilla

Structure from which a feather develops. It is composed of epidermal and dermal cells.

## Display

Behavior directed at attracting the attention of a partner. It can also be done to threaten or distract another animal.

## Distribution

Place where a species is located. It includes the area the species occupies in different seasons.

## Down Feather

A very thin and light feather, similar to silk, that birds have underneath their external plumage. Down feathers constitute the first plumage of chicks.

## Ecosystem

Community of living beings whose vital processes are interrelated and develop according to the physical factors of the same environment

## Egg

Large rounded shell, laid by a female bird,

that contains a yolk and a white. If fertilized, the egg has a tiny embryo that will develop into a chick (the chick feeds itself on the yolk and white). When ready, the chick will break the eggshell.

## Egg Tooth

Sharp calcium growth, in the shape of a tooth, that forms on the tip of a chick's bill during the embryonic phase. The chick uses the egg tooth to break its shell at birth.

## Environment

The natural conditions, such as vegetation and land, that surround animals and influence their development and behavior

## Epiphysis

Endocrine gland located below the corpus callosum in the brain. It produces a hormone that regulates sexual activity.

## Evolution

Gradual process of change that a species undergoes to adjust to the environment

## Extinct

No longer existing. Many bird species are now extinct (for example, ictiornites).

## Feather

Each unit of the covering (plumage) of a bird. The feathers are composed of a hard substance called keratin. They have a long quill, to which two blades are joined. The blades—formed by a great number of barbs, uniformly distributed—give the feather its shape and color.

## Fertilization

Union of the reproductive cells of a male and a female that will create a new individual

## Field Mark

Natural distinct feature or artificial identification of an individual of a bird species

that helps ornithologists distinguish it from other individuals of the same or other avian species

## Fledgling

Very young bird that lives in the nest where it was raised

## Fossils

Vestiges of ancient creatures of different types (vegetal or animal) on a stone substratum. Fossils can be found in the geologic strata of the Earth's surface.

## Gastric Juice

Set of fluids produced by the stomach glands of birds and other animals

## Gizzard

Muscular stomach of a bird. It is very robust, especially in granivores, and it is used to grind and soften the food by means of mechanical pressure. The food arrives at the gizzard mixed with digestive juices.

## Gland

Type of structure that is present in most multicellular living beings. It produces substances that act either inside or outside a bird's body.

## Gonad

Organ that makes male or female gametes. In birds, the testicles and ovaries are gonads.

## Granivore

Bird that feeds on seeds or grains. Many birds are granivores (for example, parrots and toucans).

## Gular Sac

Skin fragment in the shape of a sac that hangs from the lower mandible of certain birds (for example, pelicans)



## Habitat

Native or natural environment of an animal or plant

## Hatching

Cracking of the eggshell so that the bird can come out

## Histologic

Related to tissues and their study. When the anatomy of a bird is studied, the tissues that form the bird's organs are analyzed.

## Hormones

Secretion of certain glands that circulates through the body. They excite, inhibit, or regulate the activities of organs or of systems of organs.

## Horn

Made of horn or of a consistency similar to that of horn. The bill of birds is hornlike.

## Hypophysis

Internal secretion organ located at the hollow of the skull's base (called *silla turca*). It is composed of two lobes: one anterior and glandular and the other posterior and nervous. The hormones produced by the hypophysis influence growth and sexual development, among other things.

## Hypothalamus

Region of the encephalon located at the cerebral base, joined to the hypophysis by a nerve stem, in which important centers for vegetative life are found

## Incubation

The act of keeping the eggs warm so that the embryos inside can grow and hatch. Usually the chick's parents use their own bodies to warm the eggs, but some birds use sand or decomposing plants to cover them.

## Insectivore

Bird that eats insects as part of its diet

## Instinct

Innate behavior that a bird or other animal develops and that is not learned. The offspring of ducks start to swim by instinct.

## Invertebrate

Animal that lacks a spinal column. Worms, crabs, and jellyfish are examples.

## Lethargy

Sleep through which a bird can reduce its cardiac rhythm and its body temperature to save energy, especially at night and during extended periods of cold

## Malpighian Layer

Layer of epithelial cells that forms the bird's epidermis

## Mangrove Swamp

Type of ecosystem often considered a type of biome. It is composed of trees that are very tolerant to salt. These trees are found in the intertidal zone of tropical coasts. Areas with mangrove swamps include estuaries and coastal zones.

## Migration

The movement of birds from one region to another; it usually takes place in the spring and fall. It is also common among other species of animals.

## Molt

Process through which birds lose old worn feathers, replacing them with new ones

## Monogamous

Birds that mate with only one individual of the opposite sex. Many penguins have monogamous behavior.

## Morphology

Study of the form of an object or structure. For instance, the morphology of the feet of birds is an area of study.

## Nectar

Sweet and sugary secretions found in flowers that attract birds and other animals. Hummingbirds feed on nectar.

## Nidicolous

A helpless chick that depends on its parents' care after birth

## Nidifugous

A chick that can move and leave the nest as soon as it breaks its shell. In less than a day, such chicks can move agilely.

## Nocturnal

Active at night. Many birds of prey, such as owls, specialize in nocturnal hunting.

## Nutrient

Any substance obtained through diet that participates in the vital functions of a living being.

## Omnivore

Bird that has a varied diet, including animal and vegetal foods

## Pelagic

Birds that live in areas over open waters, away from the coast

## Pellet (Bolus)

Small, hard mass that some birds regurgitate (vomit). It contains parts of the food that they could not digest, such as bones, fur, feathers, and teeth.

## Pigment

Substance that colors the skin, feathers, or tissues of animals and plants

## Piscivore

Birds living in continental or oceanic waters that feed on fish

## Pollution

A consequence of human actions for natural environments. The emission of industrial gases into the atmosphere, for example, produces pollution.

## Polygamy

Reproductive relationship between one animal of one sex and several animals of the other. When one male mates with several females, it is called polygyny. Only rarely do females have multiple male reproductive partners (polyandry).

## Population

Set of individuals of the same species that live together in the same space at the same time

## Predator

Animal that hunts other animals. Birds of prey hunt other birds, mammals, and vertebrates.

## Prey

Animal hunted by another to serve as food for the latter. Animals that hunt prey are called predators.

## Protein

Organic macromolecule that is part of living beings. By including proteins in their diet, birds get the necessary amino acids to build their own organs.

## Protein Cord

Embryonic structure: each of the two filaments that sustain the yolk of the egg within the white

## Proventriculus

The first portion of the stomach, or the true

stomach, of a bird. The other portion of a bird's stomach is the gizzard.

## Rectrices

Technical term used by ornithologists to describe a bird's tail feathers

## Scale

Dermic or epidermic layer that totally or partially covers the feet of birds. They are reptilian vestiges.

## Song

Sound or series of sounds produced by a bird to demarcate its territory or to find a mate. The songs of birds can be simple or elaborate, and some are very melodic.

## Songbirds

Singing birds. Passerines include songbirds.

## Species

Set of individuals that recognize themselves as belonging to the same reproductive unit

## Sternum

Central chest bone. The sternum of flying birds has a large surface in which muscles are inserted.

## Survival

A bird's ability to face the demands of its environment and of intra- and interspecies relationships

## Swamp

Depression on the ground in which water is gathered, sometimes called a marsh. Its bottom is more or less boggy. It is the habitat of many wading birds.

## Thermal

Hot air current that rises. Many birds make use of it to gain height effortlessly.

## Theropods

Group to which carnivorous dinosaurs belong

## Training

Teaching an animal new skills. Carrier pigeons are trained.

## Tundra

Vast plains without trees in the Arctic regions of northern Asia, Europe, and North America

## Uropygial Gland

Produces an oily secretion that birds, using their bills, spread on their feathers to make them impermeable

## Vertebrate

Animals that have a spinal column, such as birds, fish, reptiles, amphibians, and mammals

## Virus

Infectious agent that depends on a living being to reproduce. Avian flu is transmitted this way.

## Vulnerable

Birds that are endangered in their natural habitats

## Yolk

Yellow part of the egg. If the egg is fertilized, a small embryo grows that will use the yolk (and white) as food.

## Young

Bird or any other animal at an early stage of life. Some young show color patterns that are very different from that of adults of the same species, which makes it difficult for predators to identify them.



# Index

## A

### adaptation

aquatic life, 5  
cellulose digestion, 57  
environment, 66  
flightless birds, 29  
flying, 8f, 12, 24-25  
foot, 20-21  
hatching, 53  
marine birds, 70, 71  
perching, 13, 79  
swimming, 68  
urban habitats, 83  
walking, 7

### adaptive radiation, 62

### aero-terrestrial habitat, 65

### African ostrich

flightless birds, 68-69  
weight, 8

### Afrotropic region, 67

### albatross

birth, 52  
classification, 64  
gliding, 35  
incubation (egg), 52  
migration routes, 59  
wings, 29

### altitudinal migration, 58

### anatomy

external features, 8-9  
internal organs, 14-15  
skeleton and musculature, 12-13  
*See also specific body parts by name, for example, wing*

### ankle, 21

### annual life cycle, 42-43

### Anseriformes (bird classification), 64, 72

### Antarctic bird: *See penguin*

### apteria, 27

### Apterygiformes (bird classification), 65, 69

### aquatic bird

classification, 64, 65  
foot adaptation, 20

physical adaptations, 5

powder down, 27

*See also marine bird*

### arboreal theory (evolution), 24

### Archaeopteryx, 10-11

### arctic tern, migration, 39, 58

### aricari, bill, 19

### artery, 15

### Asia

bird domestication, 89  
migration routes, 59

### Australasia, 67

### Australian bowerbird, courtship behavior, 46

### avian influenza A, 88

### Aztec culture, 82

## B

### bald eagle

birds of prey, 74  
talons, 21

### bar-headed goose

flight altitude, 38  
migration routes, 59

### barn swallow, migration, 58, 59, 79

### bat, wing, 24

### beak: *See bill*

### Bengal vulture, 91

### bill, 18-19

birds of prey, 74  
ducks, 73  
food filtration, 56  
identification, 9  
modern bird, 12  
parrot, 76

### binocular vision, 17

### binoculars, 84

### biodiversity

endangered species, 90-91  
geographical regions, 66-67  
running birds, 69

### biogeographic region, 66-67

### bird flu: *See avian influenza A*

### bird of prey, 74-75

pesticide poisoning, 90  
vision, 17

### birdcall,

communication, 44  
warnings, 61

### birdsong, communication, 44-45

### birdwatching, 84

### black grouse, courtship display, 31

### black swan, flight speed, 38

### blackbird, urban habitats, 87

### blood circulation: *See circulatory system*

### blue-and-white swallow, 80

### blue-footed booby, courtship behavior, 43

### bone: *See skeletal system*

### brain

Archaeopteryx, 10  
sound production, 44

### breathing: *See respiratory system*

### brown pelican, 70

### Bubo capensis, 6, 7

### Buffon, Georges-Louis Leclerc, Count de, 66

## C

### California condor, captive breeding, 90

### call: *See birdcall*

### camera, 84

### camouflage

defense strategies, 60  
molting, 32

### Campbell Island teal, 91

### canary, 88

### cannon net, 85

### Cape gannet

marine birds, 70  
migration routes, 58

### captive breeding, 90

### carnivorous bird

gizzard, 14  
*See also bird of prey*

### carpal bone, 13

### carpometacarpus, 13

### Caspian tern, migration routes, 59

### Casuariiformes (bird classification), 65, 69

### Central America, migration routes, 58

### Charadriiformes (bird classification), 64, 65, 73

### China, birds in symbolism, 82

### chough, flight altitude, 38

### circulatory system, 15

### classification, 64-65

### claw, 21

### climbing, foot adaptation, 20

### cockatoo, endangered species, 91

### collarbone

Archaeopteryx, 11  
modern bird, 12-13

### coloration, 77

flesh, 12  
seasonal changes, 32

### common loon, 70

### common spoonbill, feeding, 73

### common waxbill, 20

### communication, 44-45

### compass-and-triangulation system (migration), 59

### condor

captive breeding, 90  
dimensions, 75

### contour (feather), 26

### coracoid, 13

### courtship, 46-47

annual life cycle, 43  
black grouse, 31  
hummingbird, 37  
tail feathers 30

### covert (feather), 26

### crane

courtship behavior, 47  
migration routes, 59

### crop (esophagus), 14

### crossbill, bill, 19

### curlew sandpiper, migration routes, 59

### cursor theory (evolution), 24

## D

### Darwin, Charles, 10

### data collection, 85

### defense

camouflage: *See camouflage*  
claw, 21  
individual behavior, 60  
group behavior, 61

### diet, 56-57

adaptation of bill, 18  
courtship food, 46  
dive fishing, 71  
flight needs, 36-37  
nidicolous birds, 55  
nidifugous birds, 54  
Universe of Birds, 5

### digestive process, 14

### digestive system, 14-15

### dinosaur fossil, 10

### diurnal bird of prey, 74

*See also specific birds by name, for example, eagle*

### diving bird, 73

### dodo, 91

### domestication, 88-89

### dove, symbolism, 83

### down (feather), 26

### duck

feeding, 62, 72  
human consumption, 88  
swimming mechanism, 72

### dust bath, 27

## E

### eagle

birds of prey, 74-75  
carrying ability, 25  
feeding strategies, 57  
migration routes, 59  
speed, 39

symbolism, 82-83

talons, 20, 21

### ear

external anatomy, 9  
senses, 16

### egg, 50-51

incubation, 52

### egg tooth, 53

### Egyptian mythology, birds in symbolism, 82-83

### eider, flight speed, 38

### embryo formation, 50-51

### endangered species, 90-91

### epiphysis: *See pineal gland*

### equipment, bird watching, 84-85

### esophagus, crop, 14

### Eurasian eagle owl, 74

### Europe, migration routes, 58-59

### evolution, 10-11

flight adaptation theories, 24

### excretory system, 14

### extinction, risk, 91

*See also endangered species*

### eye, 17

### eyesight: *See vision*

## F

### falcon

bill, 19  
speed, 39  
urban habitats, 86

### falconry, 82

### false knee: *See ankle*

### farm production, 88-89

### feather

camouflage, 60  
color, 77  
external anatomy, 8-9  
flight, 26-27  
growth and molting, 32-33  
human uses, 83  
parrot, 22-23  
seasonal color changes, 32



tail function, 30  
*See also tail*

**feeding strategy**, 57

**femur**, 13

**finch**, 90

**fish, in diet**, 57

**fishing method**, 71

**flamingo**  
 bill, 19  
 food filtration system, 56  
 milk feeding, 56-57

**flapping flight**, 36-37

**flapping wing**, 12

**flight, 22-39**  
 adaptations and mechanisms, 8, 12  
 diet, 56-57  
 oxygen needs, 14  
*See also wing*

**flight pattern**, 35

**flightless bird, 68-69**  
 geographic distribution, 67  
 wings, 29

**flying animal**, 25

**foot**, 20-21  
 Archaeopteryx, 11  
 birds of prey, 75  
 external anatomy, 9  
 marine adaptations, 70  
 perching adaptations, 13, 79  
 skeleton and musculature, 13  
 zygodactyl, 77

**fossil record**  
 bones, 24  
 German findings, 10

**freshwater bird**, 72-73

**frigatebird**, reproductive season, 43

**fruit**, in diet, 57

**fulvous whistling duck**, 72-73

**furculum**: *See collarbone*

## G

**Galliformes** (bird classification), 65, 69

**geographic distribution**, 66-67  
 Passerines, 79

**Germany**, fossil findings, 10

**gizzard**, types, 14

**gliding**, 34-35

**golden plover**  
 flight endurance, 39  
 migration routes, 58

**goose**  
 domestic breeds, 88  
 flight altitude, 38  
 flight pattern, 35

**goshawk**  
 birds of prey, 74  
 courtship behavior, 46

**grain food**, 57

**granivorous bird**, gizzard, 14

**grasping mechanism**, 13  
 foot adaptation, 20

**great crested grebe**  
 courtship behavior, 46  
 lobed toes, 21

**greenfinch**, bill, 19

**grey crowned crane**, courtship behavior, 47

**group defense behavior**, 61

**growth stage**  
 nidicolous birds, 55  
 nidifugous birds, 54

**gull**  
 herring gull, 70  
 urban habitats, 86

## H

**habitat**, 65  
 destruction, 90  
 direct observation, 84  
 terrestrial, 65  
 urban habitats, 83, 86-87  
 woodpecker, 76  
*See also aquatic bird; freshwater bird; marine bird*

**hammerkop**, feeding, 73

**hatching**, 52-53  
 postnatal development, 54-55  
*See also egg*

**hatching muscle**, 53

**hawk**, bill, 74

**hearing**, 16

**heart**, 15

**Heligoland trap**, 85

**heron**  
 bill, 19  
 feeding, 73  
 imperial heron, 27  
 tricolored heron, 20-21  
 white heron, 4, 5

**herring gull**, 70

**hoatzin**, 64, 66

**Holarctica**, 67

**human anatomy**  
 comparison, 10  
 leg, 21  
 vision compared, 17

**human interaction**, 82-83  
 domestication, 88-89

**humerus**, 13

**hummingbird**  
 bill, 19  
 courtship behavior, 37  
 external anatomy, 12  
 flight mechanism, 37  
 internal organs, 14-15  
 migration routes, 58  
 Passeriformes, 78  
 ruby-throated hummingbird, 58, 67  
 tongue, 57  
 wine-throated hummingbird, 8

**hunting bird**: *See bird of prey*

**hyacinth macaw**, 90

**hypophysis**, 42

## I

**ibis**, 73

**identification**, 9

captured birds, 85  
*See also classification*

**imperial heron**, 27

**imperial shag**, 70

**incubation** (egg), 52

**incubator bird**, 54

**India**  
 domestication, 89  
 representation of god, 82

**Indian vulture**, 91

**Indomalaya region**, 67

**insectivore**, 57

## J

**jaw**, 19

**jawbone**, Archaeopteryx, 10

**Jurassic Period**, Archaeopteryx, 10-11

## K

**keratin**, 27

**kiwi**, 69

**knee**, 21

## L

**latitudinal migration**, 58

**laying (egg)**: *See egg*

**Leclerc**, Georges-Louis, Count de Buffon, 66

**leg**, 20-21  
 grasping mechanism, 13  
 muscles, 13

**lek ritual**, 46

**lesser spotted eagle**, migration routes, 59

**life cycle**, annual, 42-43

**little masked weaver**, nest building, 49

**longitudinal migration**, 58

**loon**: *See common loon*

**lung**, 15

**lyrebird**, 78

## M

**Malaysian cockatoo**, 91

**mangrove finch**, 90

**marine bird**, 65, 70-71  
 gliding, 35  
*See also aquatic bird*

**marine glider**, 34

**marking** (captured bird), 85

**Mayan culture**, 82

**meat, in diet**, 57

**metabolism**, 9

**Mexico**, domestication, 89

**migration**, 58-59  
 arctic tern, 39  
 Passerines, 79  
 Universe of Birds, 4-5

**mimicry of sound**, 76, 78, 79

**mist net**, 85

**molting**, 32-33

**monocular vision**, 17

**monogamy**, 47

**musculature**, 12-13

**mythology**, 82-83

## N

**nasal marker**, 85

**Nearctic region**, 66

**neck marker**, 85

**nectar**, 57

**Neotropic region**, 66

**nest**  
 construction, 49  
 types, 48

**nesting**, 5  
 partridge, 41

**net**, types, 85

**nidicolous bird**, 54, 55

**nidifugous bird**, 54

**nocturnal bird**, 74  
*See also owl*

**North America**  
 indigenous peoples, 83  
 migration routes, 58

**northern harrier**, courtship behavior, 46

**northern wheater**, migration routes, 59

**nostril**, 71

**nuptial display**: *See courtship*

## O

**Oceania**, 66

**Origin of Species, The** (Darwin), 10

**ornithology**, 84-85

**ostrich**  
 African ostrich, 8, 68-69  
 distribution, 67

**ovenbird**, 79

**owl**  
 birds of prey, 74  
*Bubo capensis*, 6-7  
 diet, 57  
 urban habitats, 86

**oxygen**, consumption, 14

## P

**Palaearctic region**, 67

**parental care** (evolution), 24

**parrot**  
 feathers, 22-23  
 flight speed, 38  
 talking birds, 76

**partridge**  
 nesting, 41  
 red-legged partridge, 54

**passerine (Passeriformes)**, 78-79  
 external anatomy, 9



classification, 64, 65

**pelican**

- brown pelican, 70
- flight, 34-35
- resting pose, 42-43

**pelvis**, Archaeopteryx, 11

**penguin**

- external anatomy, 8
- feather coverage, 27
- incubation (egg), 52
- mating behavior, 47
- molting, 32
- observation, 85
- swimming adaptations, 68
- swimming speed, 39

**perching**

- foot adaptation, 20
- skeleton and musculature, 13
- See also **passerine**

**peregrine falcon**

- migration routes, 58
- pesticide poisoning, 90
- speed, 39

**pesticide**, poisoning, 90

**pheasant**, flight speed, 38

**pigeon**

- incubation (egg), 52
- messenger service, 88
- milk feeding, 56-57
- skeleton, 10
- urban habitats, 83, 86
- wing mobility, 11

**pineal gland**, 42

**plumage**: See **feather**

**pneumatic bone**, 13

**Podicipediformes** (bird classification), 65, 73

**polyandry**, 47

**polygamy**, 47

**polygyny**, 47

**population control**, urban habitats, 87

**postnatal development**, 54-55

**poultry farming**, 88-89

**powder down**, 27

**preening**, 27

**ptarmigan**, molting, 32

**pterodactyl**, wing, 24

**pterylae**, 27

**puffin**, 66

**pygostyle**, 11, 13

See also **tail**

## Q-R

**quetzal**, 76, 82

**radius**, 13

**ratite**, 69

**raven**, 19, 78

**religion**, birds as symbols, 82-83

**reproduction**

- courtship: See **courtship**
- egg, 50-51
- hypophysis, 42

**reptile**, bird evolution, 24

**respiratory system**, 15

**rhea**, 69

**ribs**, Archaeopteryx, 11

**ringed bird**, 85

**robin**, 80-81, 87

**rockhopper penguin**, 68

**royal cinchlo**, 90

**royal eagle**, speed, 39

**royal swift**, flight speed, 38

**ruby-throated hummingbird**

- bird distribution, 67
- migration routes, 58

**ruff**, courtship behavior, 43

**running bird**

- flightless birds, 69
- foot adaptation, 20

**running theory** (evolution), 24

**Ruppell's griffon vulture**, 39

## S

**salt gland**, 71

**scale** (bird skin)

feet, 21

types, 25

**Scots dumpy rooster**, spurs, 21

**seagull**, urban habitats, 86

**seasonal color change**, 32

**seed food**, 57

**seizing**: See **grasping mechanism**

**sense**, 16-17

- external anatomy, 9

**shoebill**, feeding, 73

**Siberian crane**, migration routes, 59

**sight**: See **vision**

**skeletal system**, 12-13

- Archaeopteryx, 11
- legs, 21
- running birds, 69

See also *under specific topic, for example toe*

**skull**

- Archaeopteryx, 10
- modern bird, 12

**sleep regulation**, 42

**smell**, 16

**snow bunting**, migration routes, 58

**snowy egret**, courtship behavior, 46

**song**: See **birdsong**

**songbird**

- classification, 64, 65
- distinction, 9
- Passeriformes, 78-79

**sound production**

- recording, 84
- songbirds, 78, 79
- talking birds, 76

**South America**

- domestication, 89
- migration routes, 58
- species diversity, 66

**sparrow**

- urban habitats, 83, 87
- white-throated sparrow, 8-9

**sparrowhawk**, 74

**species classification**: See **classification**

**speed**, 38-39

**spin tailed swift**, speed, 39

**spine**

- Archaeopteryx, 10
- bones, 13

**spoonbill**: See **common spoonbill**

**spur**, 21

**stability**, 8

**starling**

- defense formation, 61
- flight speed, 38

**sternum**, modern bird, 12-13

**stork**

- feeding, 73
- flight speed, 38
- migration routes, 58
- urban habitats, 86

**Struthioniformes** (bird classification), 65, 69

**swallow**, 78, 79, 80

**swan**

- flight altitude, 38
- migration routes, 59

**swift**, urban habitats, 86

**symbolism**, 83

**syrix**, sound production, 44, 79

## T

**tail**, 30-31

- Archaeopteryx, 11
- pygostyle, 11, 13

**tail feather**, 8

**talon**, Archaeopteryx, 11

**tarsometatarsus**, 13

- Archaeopteryx, 11

**terrestrial animal**, speed of movement, 39

**terrestrial glider**, 34, 35

**terrestrial habitat**, 65

**territorial demarcation**, 45

**theropod**, 10

**thigh**, 21

**tibia**, 13, 21

**tibiotarsus**, 21

**toe**

- anatomy, 9, 13
- Archaeopteryx, 11

- types, 20

**tongue**

- hummingbird, 57
- internal organs, 15, 16

**toucan**

- bill, 19
- colorful birds, 76

**touch**, 16

**Triassic Period**, theropodan reptile, 10

**tricolored heron**, 20-21

**turkey**, 88

## U-V

**ulna**, 13

**urban habitat**, 83, 86-87

**vein**, 15

**velociraptor**, 10

**vibrissae**, 27

**vision**, 16, 17

**vulture**

- endangered species, 91
- feeding practices, 75
- flight altitude, 39

## W

**wading bird**, 73

**walking bird**, adaptations, 7, 20

**wandering albatross**

- migration routes, 59
- distances, 40
- wings, 29

**water absorption**, 14

**waxbill**: See **common waxbill**

**white heron**, 4, 5

**white stork**, migration routes, 58

**white-throated sparrow**, 8-9

**wide bill**, 79

**wine-throated hummingbird**, 8

**wing**, 28-29

- birds of prey, 75
- feathers, 9
- gliding, 34-35
- markers, 85
- skeleton and musculature, 12-13
- types and evolution, 24

**wing loading**, 29

**winglet**, 34

**wingspan**, 29

**woodpecker**, 76

**wrist joint**, Archaeopteryx, 11

## Y-Z

**yellow-crested cockatoo**, 91

**zone-tailed hawk**, 74-75





## BIRDS

Britannica Illustrated  
Science Library



ENCYCLOPEDIA  
Britannica<sup>®</sup>